

Managing the spread of *Pseudogymnoascus Destructans* and conserving Bats threatened by White-Nose Syndrome in North America

Jeremy T. H. Coleman¹, Jonathan D. Reichard¹, Richard Geboy², Christopher Servheen^{3*},
Christina Kocer¹, and Mike Armstrong⁴

U.S. Fish and Wildlife Service: ¹Hadley, Massachusetts; ²Bloomington, Indiana; ³Missoula, Montana; ⁴Frankfort, Kentucky

Bats in North America face numerous challenges, including habitat loss, climate change, environmental contamination, and energy development, but none is as immediate a threat to multiple species as white-nose syndrome, a fungal disease that is responsible for unparalleled rapid declines in bat populations. White-nose syndrome (WNS) was unknown to science before discovery in New York in January 2007 (Blehert *et al.* 2009). Since that time the disease has spread to at least 26 states and five Canadian provinces, and the fungus causing the disease, *Pseudogymnoascus destructans* (*Pd*, Lorch *et al.* 2011), has been documented in three additional states: Minnesota, Mississippi, and Oklahoma (see current map: www.WhiteNoseSyndrome.org). Genetic analyses indicate the fungus is likely not native to North America and that human activity may have led to its introduction. *P. destructans* may have originated in Europe where the fungus has been detected on 13 bat species, 11 of which have been confirmed with WNS based on our current case definitions. There have been no documented deaths of European bats from WNS, however.

From what appears to be a single point of introduction to North America, *Pd* has spread rapidly, propagated largely by bat-to-bat or bat-to-environment-to-bat contact. Human-mediated transmission also may have contributed to the spread based on the potential for the fungus to be carried between caves on clothing or personal gear (USGS 2009), the ability of the fungus to remain viable for long

durations in the absence of bat hosts (Lorch *et al.* 2013, Hoyt *et al.* 2014), and conspicuous longer distance jumps to heavily visited caves before detection at large bat hibernacula close by. The fungus infects torpid bats resulting in physiological and behavioral impacts, often leading to mortality. Population declines exceeding 90% have been documented in affected hibernacula during winter surveys (Turner *et al.* 2011), and corroborated by estimates of colony size and activity during summer. The international response to the disease has been rapid, and has resulted in considerable advances in our understanding of the pathogen and the disease to the extent that we are beginning to see progress in the development of potential tools and strategies designed to better manage *Pd* and the impacts of WNS on bats.

In the United States, WNS research, management, and conservation activities are coordinated through a common collaborative strategy outlined in: *A National Plan for Assisting States, Federal Agencies and Tribes in Managing White-Nose Syndrome in Bats* (USFWS 2011). The U.S. national plan is mirrored in Canada by a sister plan: *A National Plan to Manage White-Nose Syndrome in Bats in Canada* (CWHC 2015). Despite the advances made in our understanding of the mechanics of this epizootic, including pathology, transmission, dynamics, and ecological requirements of *Pd*, the best tools currently available are still those

Table 1. Bat species on which white-nose syndrome (WNS) has been confirmed based on approved case definitions, or on which at least one individual has tested positive for *Pseudogymnoascus destructans* (*Pd*) using molecular identification techniques

	<u>Positive for WNS</u>	<u>Pd Present</u>
North America		
	Big brown bat (<i>Eptesicus fuscus</i>) ¹	Eastern red bat (<i>Lasiurus borealis</i>) ²
	Eastern small-footed bat (<i>Myotis leibii</i>)	Southeastern bat (<i>M. austroriparius</i>)
	Gray bat (<i>M. grisescens</i>) * endangered	Silver-haired bat (<i>Lasionycteris noctivagans</i>) ²
	Indiana bat (<i>M. sodalis</i>) * endangered	Rafinesque's big-eared bat (<i>Corynorhinus rafinesquii</i>) ²
	Little brown bat (<i>M. lucifugus</i>) ¹	Virginia Big-Eared Bat (<i>C. townsendii virginianus</i>) * endangered
	Northern long-eared bat (<i>M. septentrionalis</i>) ¹ * threatened	
	Tri-colored bat (<i>Perimyotis subflavus</i>) ¹	
Europe		
	Greater mouse-eared bat (<i>M. myotis</i>) ³	Whiskered bat (<i>M. mystacinus</i>) ⁵
	Daubenton's bat (<i>M. daubentonii</i>) ⁴	Lesser mouse-eared bat (<i>M. oxygnathus</i>) ⁶
	Bechstein's bat (<i>M. bechsteinii</i>) ⁴	
	Natterer's bat (<i>M. nattereri</i>) ⁴	
	Brandt's bat (<i>M. brandtii</i>) ⁴	
	Geoffroy's bat (<i>M. emarginatus</i>) ⁴	
	Pond bat (<i>M. dasycneme</i>) ⁴	
	Northern bat (<i>E. nilssonii</i>) ⁴	
	Barbastelle (<i>Barbastellus barbastellus</i>) ⁴	
	Brown long-eared bat (<i>Plecotus auritus</i>) ⁴	
	Lesser horseshoe bat (<i>Rhinolophus hipposideros</i>) ⁴	
1 - Blehert et al. 2009; 2 - Bernard et al. 2015; 3 - Pikula et al. 2012; 4 - Zupal et al. 2014; 5 - Martinkova et al. 2010; 6 - Wibbelt et al. 2010		

that reduce the risks of human-mediated transmission of the pathogen and reduce disturbance to bats during hibernation.

In North America, seven bat species have been confirmed with WNS, and five additional species have been detected carrying *Pd* (Table 1). *P. destructans* is a psychrophilic (cold-loving) fungus that grows

best in cool, wet conditions (growth range between ~ 1 and 19°C, Verant et al. 2012). Such conditions are common for caves and mines across the continent (Perry 2013), and hibernating bats in every state may be found using these same cool and humid conditions. Like most fungi, *Pd* produces spores and hyphae (filamentous

Table 2. Evidence and supporting publications or sources providing evidence that human actions may contribute to the spread of *Pseudogymnoascus destructans* (*Pd*) and affect survival of bats affected by white-nose syndrome (WNS).

<u>Evidence</u>	<u>Supporting publications and reports</u>
<i>Pd</i> can persist and grow in the absence of bats.	Lindner et al., 2011; Lorch et al., 2012; Lorch et al., 2013; Puechmaille et al., 2011a; Reynolds et al. 2015
Spores of <i>Pd</i> can remain viable outside of subterranean environments.	Puechmaille et al., 2011b; Hoyt et al. 2014
Spores of <i>Pd</i> can cling to clothing and footwear and can be inadvertently transported out of contaminated sites.	J. Okoniewski, unpublished data; H. Barton, unpublished data; USGS - National Wildlife Health Center, unpublished data
<i>Pd</i> may be present on bats or in bat roosts without being visibly detectable.	Dobony et al., 2011; Langwig et al., 2015
Spread of <i>Pd</i> may be slowed by geographic or biological barriers to bat movements that may not be barriers to human movement.	Miller-Butterworth et al., 2014
Repeated and/or prolonged human disturbance during hibernation is detrimental to bats, especially bats already stressed by WNS.	Boyles and Willis, 2009; McCracken, 1989; Mohr, 1972; Reeder et al., 2012; Thomas, 1995; Tuttle, 1979
If done correctly, current decontamination procedures have a high probability of significantly reducing the risk of spreading viable <i>Pd</i> .	Shelley et al., 2013

vegetative growth), and is capable of asexual reproduction through spores and hyphal fragmentation. The spores can persist for extended periods in suboptimal conditions, including wide temperature and humidity ranges (Lorch *et al.* 2013). Fungal material can adhere to substrate, fur, hair, clothing, shoes, etc. (collectively referred to as “fomites”), and thus, can be moved large distances by bats, humans, or other carriers. As a consequence, human visitation to caves and mines increases the risk of contributing to the spread of WNS to uncontaminated areas through inadvertent transport of *Pd*. The containment strategies developed for WNS include: 1) cleaning and treating (hereafter, decontamination) of gear, clothing, footwear, and any other surfaces that might have been exposed to *Pd* and may subsequently come in contact with hibernating bats or their habitats, and 2) the reduction or prevention of opportunities for people to encounter, transport, and transfer *Pd* spores or

vegetative structures to bats or naïve environments.

The U.S. Fish and Wildlife Service developed guidance for decontamination procedures (Decontamination Protocol, 2008) as well as recommendations for regulation or restriction of human activities likely to pose a risk for spreading *Pd* (Cave Advisory, 2009). These documents were developed in consultation with state and federal partners that would later join together formally under the collaborative national response plan. The national plan provided further support for the importance of containment strategies for managing infectious agents, and describes the ways in which universal precautions are implemented to reduce incidence of disease by preventing infections and breaking chains of transmission. The Decontamination Protocol has been revised regularly in response to research results that further our understanding of the compounds and procedures known to effec-

tively kill spores of *Pd* (e.g., Shelley *et al.* 2013), and to include a wider variety of applications. The Cave Advisory has also been reviewed regularly, and has been the focus of considerable discussion within the WNS response community, but had not been formally revised until recently (revision under final review). While some have questioned the risks associated with human-mediated spread of *Pd*, the weight of scientific evidence in support of these protocols and recommendations continues to grow (Table 2). Additionally, members of the WNS Steering Committee have agreed to endorse the most recent versions of the two documents as national guidance, so they now represent an agreed upon set of recommendations from state, federal, and tribal

agencies engaged in the national response. Updated versions of the National Decontamination Protocol and National Recommendations for Managing Access to Subterranean Bat Roosts (replacing the Cave Advisory) will be released in coming months.

Dedication and commitment of the many individuals, institutions, and government agencies who have engaged in this issue and lent their support has produced both great progress in our understanding of WNS and ability to respond to it, and significant contributions to broader scientific advancement. Healthy bat populations contribute to strong ecosystems and provide beneficial ecosystem services, thus our efforts to control and reduce the impacts of WNS on North American bats is of the highest priority.

References

- Bernard, R.F., J.T. Foster, E.V. Willcox, K.L. Parise, and G.F. McCracken. 2015. Molecular detection of the causative agent of white-nose syndrome on Rafinesque's big-eared bats (*Corynorhinus rafinesquii*) and two species of migratory bats in the southeastern USA. *Journal of Wildlife Diseases*. 51:519-522.
- Blehert, D. S., A. C. Hicks, M. Behr, C. U. Meteyer, B. M. Berlowski-Zier, E. L. Buckles, J. T. H. Coleman, S. R. Darling, A. Gargas, R. Niver, J. C. Okoniewski, R. J. Rudd, & W. B. Stone. 2009. Bat white-nose syndrome: an emerging fungal pathogen? *Science* **323**:227.
- Boyles, J.G., and C.K.R. Willis. 2010. Could localized warm areas inside cold caves reduce mortality of hibernating bats affected white-nose syndrome? *Frontiers in Ecology and the Environment* **8**: 92-98.
- Canadian Wildlife Health Cooperative (CWHC). 2015. A National Plan to Manage White-Nose Syndrome in Bats in Canada. Saskatoon, Saskatchewan.
- <<http://www.cwhc-rcsf.ca/docs/BatWhiteNoseSyndrome-NationalPlan.pdf>> Accessed September 10, 2015.
- Dobony, C. A., A. C. Hicks, K. E. Langwig, R. I. von Linden, J. C. Okoniewski, & R. E. Rainbolt. 2011. Little brown myotis persist despite exposure to white-nose syndrome. *Journal of Fish and Wildlife Management* **2**:190–195.
- Fisher, M. C., D. A. Henk, C. J. Briggs, J. S. Brownstein, L. C. Madoff, S. L. McCraw, & S. J. Gurr. 2012. Emerging fungal threats to animal, plant and ecosystem health. *Nature* **484**:186–194.
- Hoyt, J. R., K. E. Langwig, J. Okoniewski, W. F. Frick, W. B. Stone, & A. M. Kilpatrick. 2014. Long-Term persistence of *Pseudogymnoascus destructans*, the causative agent of white-nose syndrome, in the absence of bats. *EcoHealth* **16**:12-202. DOI: 10.1007/s10393-014-0981-4.
- Langwig, K.E., W.F. Frick, R. Reynolds, et al. 2015. Hibernation drives seasonal dynamics of a fungal disease. *Proceedings of the Royal Society B*. 2015 282 20142335; DOI: 10.1098/rspb.2014.2335.
- Lindner, D.L., A. Gargas, J.M. Lorch, M.T. Banik, J. Glaeser, T.H. Kunz, and D.S. Blehert. 2011. DNA-based detection of the fungal pathogen *Geomyces destructans* in soils from bat hibernacula. *Mycologia* **103**:241-246.
- Lorch, J. M., C. U. Meteyer, M. J. Behr, J. G. Boyles, P. M. Cryan, A. C. Hicks, A. E. Ballmann, J. T. H. Coleman, D. N. Redell, D. M. Reeder, & D. S. Blehert. 2011. Experimental infection of bats with *Geomyces destructans* causes white-nose syndrome. *Nature* **480**:376-378.

- Lorch, J.M., D.L. Lindner, A. Gargas, L.K. Muller, A.M. Minnis, and D.S. Blehert. 2012. A culture-based survey of fungi in soil from bat hibernacula in the eastern United States and its implications for detection of *Geomyces destructans*, the causal agent of bat white-nose syndrome. *Mycologia*. doi:10.3852/12-207
- Lorch, J. M., L. K. Muller, R. E. Russell, M. O'Connor, D. L. Lindner, & D. S. Blehert. 2013. Distribution and environmental persistence of the causative agent of white-nose syndrome, *Geomyces destructans*, in bat hibernacula of the eastern United States. *Applied and Environmental Microbiology* **79**:1293-1301.
- Martínková, N., P. Bačkor, T. Bartonička, P. Blažková, J. Červený, *et al.* 2010. Increasing incidence of *Geomyces destructans* fungus in bats from the Czech Republic and Slovakia. *PLoS One* **5**:e13853.
- McCracken, G. 1989. Cave conservation: special problems for bats. *The NSS Bulletin* **51**: 47-5.
- Miller-Butterworth, C. M., M. J. Vonhof, J. Rosenstern, G. G. Turner, & A. L. Russel. 2014. Genetic Structure of Little Brown Bats (*Myotis lucifugus*) Corresponds With Spread of White-Nose Syndrome Among Hibernacula. *Journal of Heredity* **105**:354-364.
- Mohr, C.E. 1972. The status of threatened species of cave dwelling bats. *Bulletin of the National Speleological Society* **34**: 33-47.
- Perry, R. W. 2013. A review of factors affecting cave climates for hibernating bats in temperate North America. *Environmental Reviews* **21**:28-39.
- Pikula, J., H. Bandouchova, L. Novotny, C. U. Meteyer, J. Zupal, *et al.* 2012. Histopathology confirms white-nose syndrome in bats in Europe. *Journal of Wildlife Diseases* **48**:207-211.
- Puechmaille S.J., G. Wibbelt, V. Korn, H. Fuller, F. Forget, *et al.* 2011a. Pan-european distribution of white-nose syndrome fungus (*Geomyces destructans*) not associated with mass mortality. *PLoS ONE* **6**: e19167. doi: 10.1371/journal.pone.0019167
- Puechmaille, S.J., H. Fuller, and E.C. Teeling. 2011b. Effect of sample preservation methods on the viability of *Geomyces destructans*, the fungus associated with white-nose syndrome in bats. *Acta Chiropterologica* **13**:217-221.
- Reeder, D. M., C. L. Frank, G. G. Turner, C. U. Meteyer, A. Kurta, *et al.* 2012. Frequent Arousal from Hibernation Linked to Severity of Infection and Mortality in Bats with White-Nose Syndrome. *PLoS One* **7**: e38920
- Reynolds, H.T., T. Ingersoll, and H.A. Barton. 2015. Modeling the environmental growth of *Pseudogymnoascus destructans* and its impact on the white-nose syndrome epidemic. *Journal of Wildlife Diseases* **51**: 318-331.
- Shelley, V., S. Kaiser, E. Shelley, T. Williams, M. Kramer, K. Haman, K. Keel, and H.A. Barton. 2013. Evaluation of strategies for the decontamination of equipment for *Geomyces destructans*, the causative agent of white-nose syndrome. *Journal of Cave and Karst Studies* **75**: 1-10. doi: 10.4311/2011LSC0249
- Thomas, D.W. 1995. Hibernating bats are sensitive to nontactile human disturbance. *Journal of Mammalogy* **76**: 940-946.
- Turner, G. G., D. M. Reeder, & J. T. H. Coleman. 2011. A Five-year Assessment of Mortality and Geographic Spread of White-nose Syndrome in North American Bats and a Look to the Future *Bat Research News* **52**:13-27.
- Tuttle, M.D. 1979. Status, causes of decline, and management of endangered gray bats. *The Journal of Wildlife Management* **43**: 1-17.
- United States Fish and Wildlife Service (USFWS). 2011. A national plan for assisting states, federal agencies and tribes in managing White-Nose Syndrome in Bats. Hadley, Massachusetts. <<http://www.whitenosesyndrome.org/national-plan/white-nose-syndrome-national-plan>> Accessed August 17, 2015
- United States Geological Survey National Wildlife Health Center (USGS). 2009. Update on white-nose syndrome. *Wildlife Health Bulletin* 2009-03. 11 December 2009. <http://www.nwhc.usgs.gov/publications/wildlife_health_bulletins/WHB_2009-03_WNS.pdf> Accessed 1 October 2014.
- Verant, M. L., J. G. Boyles, W. Waldrep JR., G. Wibbelt, & D. S. Blehert. 2012. Temperature-dependent growth of *Geomyces destructans*, the fungus that causes bat white-nose syndrome.