# Western Kentucky University TopSCHOLAR®

**Nursing Faculty Publications** 

School of Nursing

1996

# Multidimensional Factors Affecting Well-Being: A PNI Based Model for Therapeutic Nursing Intervention.

Mary P. Bennett
Western Kentucky University, mary.bennett@wku.edu

Follow this and additional works at: http://digitalcommons.wku.edu/nurs\_fac\_pub

Part of the <u>Analytical</u>, <u>Diagnostic and Therapeutic Techniques and Equipment Commons</u>, <u>Nursing Commons</u>, and the <u>Psychological Phenomena and Processes Commons</u>

### Recommended Repository Citation

Bennett, Mary P., "Multidimensional Factors Affecting Well-Being: A PNI Based Model for Therapeutic Nursing Intervention." (1996). *Nursing Faculty Publications*. Paper 46. http://digitalcommons.wku.edu/nurs\_fac\_pub/46

This Other is brought to you for free and open access by TopSCHOLAR\*. It has been accepted for inclusion in Nursing Faculty Publications by an authorized administrator of TopSCHOLAR\*. For more information, please contact todd.seguin@wku.edu.

Multidimensional Factors Affecting Well-Being: A PNI

Based Model for Therapeutic Nursing Intervention.

Mary P. Bennett

**RUSH School of Nursing** 

# Introduction:

This paper presents a nursing model, gives some examples of how it could be implemented, and proposes some outcome objectives for its evaluation. Based partially on psychoneuroimmunology theory (PNI), this model was designed to help meet my own needs for a testable nursing model, and for all the nursing students who ever struggled to understand nursing theory. Perhaps those of you who can't quite relate to black holes and spiraling energy fields may find it useful in your search for the true meaning of nursing.

What is nursing? What do nurses do? Each of us knows inherently what we are and what we do, yet when asked to define it for the public (or our instructors) we find ourselves speechless. It's like trying to tell someone how to tie a shoe. The knowledge and ability are there on a level that is not just cognitive or verbal, but lies in the very connection between your mind and your fingers.

Yet, we meet with people who really do not know or understand what nurses do, let alone understand how the different educational levels relate to one another. A hospital controller once told me about a nurse practitioner who recently went into private practice, after working in collaboration with a local physician for a couple years. The controller wanted to know if it was legal, let alone ethical, for a nurse practitioner to go into practice for herself, "especially after Dr. X just finished training her". This scenario highlights how poor a job we have done in educating others about our various levels of nursing practice, when even those in the health care arena are confused.

So how can we define nursing in a way that truly encompasses everything that nurses do and all the various nursing roles available? A broad definition is clearly needed, but one that separates the role of nursing from other health care professionals. For the purpose of this theory, nursing can be defined as the art of using therapeutic nursing interventions to improve well-being. This leads to the definition of therapeutic nursing intervention. According to Buchanan (1994):

Therapeutic Nursing Interventions are nursing actions and interactions, executed as part of the nursing process, with or for individuals and families, that are directed at influencing a measurable change in health status and quality of life. Interventions are directed at: treating acute illness; alleviating illness related symptoms; promoting human adaptation to acute illness during acute and transitional phases of recovery; rehabilitation during chronic illness and disability over the life span; preventing illness; and supporting individuals and families during terminal or end of life transitions (p. 190).

Buchanan's definition requires the interventions to be part of the nursing process, which helps separate nursing interventions (at least conceptually) from those that interventions that might be applied by other health care professionals under their scope of practice. The nursing process may be viewed by some as being somewhat outdated, but even when not followed in a formal fashion, this process does provide a basic underlying framework to help explain how the nurse assesses the patient situation, selects nursing interventions to apply, and monitors patient outcomes to determine the success of the intervention. Buchanan's definition is somewhat biased towards illness care, but the definition also includes prevention of illness and supporting people during the end of life, when neither prevention nor cure of the illness is a primary focus. A more problematic issue with Buchanan's definition is the requirement that nursing interventions be directed towards producing a "measurable change". While this requirement is laudable from the standpoint of a quantitative researcher, it may be that not all beneficial outcomes can be measured in an quantitative fashion. An updated definition of therapeutic nursing intervention could be stated as:

Therapeutic Nursing Interventions: Actions performed by nurses within their legal scope of practice, educational preparation and skill level, to improve well-being in individuals, families and communities. These interventions may be directed at prevention or treatment of illness or injury, prevention of complications, and providing support to patients and their families during adaptation to changes in wellbeing and at the end of life.

However, the number and type of interventions which are available to the nurse, what type of clients are being treated, and what kind of environments are being practiced in frequently vary based upon the nurses' educational preparation. While basic nursing practice involves the assessment, diagnosis and treatment of basic human health needs in stable environments with adequate supervision, advanced nursing practice is quite different. Advanced nursing practice involves the independent assessment, diagnosis and treatment of a variety of human wellness needs in a multiplicity of environments, both highly structured and non-structured. The advanced practice role also includes the following actions: collaboration with other health care professionals; mentoring and educating others in the nursing profession; research to improve nursing practice; and political activism to increase public access to needed health care. In addition, doctorally prepared advanced practice nurses have the theoretical and research experience needed to design and test new nursing interventions, in order to expand the scope of nursing knowledge and practice.

To help outline roles and skills associated with different nursing practice levels, see table 1. The various tasks and skill levels noted are intended to be general guidelines only. We all know experienced Associate degree RN's who are capable of performing many of the tasks and skills listed at the B.S.N. level, and there are many nurses educated at the doctoral level who do not perform all of the skills and tasks listed in the D.N.S./PhD column. Each of us has to decide which area of nursing practice we are going to focus our energies on, instead of continuing to be everything to everyone. In general, nursing education programs should try to educate graduates to meet the basic requirements of each level, while making additional courses available to help students develop specialized skills.

Basic	Through	Advanced	Practice	Nursing	Functions
-------	---------	----------	----------	---------	-----------

Entry Level RN	B.S.N.	M.S.N.
Roles:  1. Basic nursing    assessment, diagnosis    and treatment of    human health needs    in stable	Roles: 1. Nursing assessment, diagnosis and treatment of human health needs in fluid environments, under the supervision of APN or MD's	Roles: 1. Advanced assessment, diagnos human health needs in primar environments 2. Independent practice/collabor MD's,/OD's, depending upon p
environments, under the supervision of B.S.N. and/or	<ol> <li>Support of physiological needs</li> <li>Support of mental health needs -touch, social contact,</li> </ol>	regulations 3. Prescribes and supervises primactivities for clients
advanced practice nurse (APN) 2. Support of basic	family relationships and human caring  4. Communication and referral	Designs and implements comm programs to encourage health mental behavior changes in a g
physiological needs 3. Support of basic mental health needs - social contact, touch,	within the larger health care system  5. Various medical functions as prescribed by MD, OD, NP, or written protocols	<ul> <li>5. Communicates and refers clier allied health professionals as n</li> <li>6. Works with multiple health car funding systems in order to propossible for his/her clients</li> </ul>
human caring  1. Communication and referral within the employing organization	Coordination of multiple health care services within the health care environment     Various supervisory and administration tasks	7. Various administration and support hospital and independent  8. Coordination of patient education.  Skills:
5 . Various medical functions as prescribed by MD's or NP's	8. Various patient education tasks  Skills:  1. Nursing assessment, planning,	Advanced assessment, diagnost technical, pharmacological, coeducational, communication, a skills
6. Basic coordination of health care services within the place of employment	technical, coordination, basic educational, and supervisory skills  2. Verbal and written	Verbal communication skills to and small groups, students, pe administrators in nursing and o     Written communication skills s
Skills:  1. Basic nursing assessment, planning.	communication skills to work with both individuals and small groups 3. Computer/business skills for	disseminate nursing knowledg student/patient educational m  4. Ability to work with multiple corretrieval systems to access hear

- assessment, planning, technical and coordination skills
- 2. Basic verbal and written communication skills.
- 3. Basic computer skills for charting, database and order entry systems
- 4. Basic social skills to work with patients/ families, and as part of the nursing team

- 3. Computer/business skills for charting, scheduling, order-entry, and unit budgeting
- 4. Social and leadership skills to work within small groups and function as entry level nursing administrator

#### **Professional Maintenance**

- 1. Member of state nurses association and active at the local level
- 2. Works with local legislators to influence the local health care environment

- osis and treatment of ry care
- ration with practice and state
- mary health care
- munity health ny physical and given population
- ents to MD/OD and needed
- are providers and rovide the best care
- pervisory tasks in nt clinic sites
- ation and entry level
- ostic, planning, ordination. and supervisory
- o work with large eers and other professions
- sufficient to ge and develop materials
- computer data base retrieval systems to access health care information and research plus basic business and educational computer skills
- 5. Social and leadership skills to work within large and small groups, function as a nursing administrator and/ or educator, and work with leaders from nursing and other professions

#### Professional Maintenance

- 1. Member of one or more professional nurses organizations and active at the state level in at least one professional nursing organization
- 2. Works with local and state legislators to influence the health care environment
- 3. Yearly continuing education
- 4. Certification in specialty area

#### Roles:

D.N.S./Ph.D.

- 1. Advanced assessment, diagnosis and treatment of human health needs in primary care environments
- 2. Independent nurse practitioner, educator, nurse researcher, and/or administrator
- 3. Performs various functions in the health care community Including: overall community resource planning, overseeing community research, and working with area business leaders and legislators to improve the health of the community
- 4. Curriculum design to teach student nurses needed skills to meet basic community needs
- 5. Prescribes and supervises primary health care for clients
- 6. Designs and implements community health programs to encourage healthy physical and mental behavior changes in a given population
- 7. Communications and refers clients to MD, OD, PT, and other health care professionals
- 8. Works with multiple health care providers and funding systems in order to provide the best care possible for his/her community
- 9. Various administration and clinical research tasks in hospitals, independent clinics, and educational settings

#### Skills:

- 1. Advanced assessment, diagnostic, planning, technical, pharmacological, coordination, education, communication and supervisory
- 2. Verbal communication skills to work with large and small groups, students, peers, researchers and administrators in nursing and other professions
- 3. Written communication skills sufficient to disseminate nursing knowledge and research through presentations and publications
- 4. Advanced computer skills to access multiple database systems, Internet communication systems, and utilize various statistical and presentation packages as needed to advance the profession and science of nursing
- 5. Social and leadership skills to work within large and small groups, function as a nursing administrator and/or educator, and work with peers, administrators, and researchers from nursing and other professions
- 6. Basic and applied clinical research skills including data analysis and research methodology


#### **Professional Maintenance**

- 1. Charting a course for the future of the nursing profession itself, given changing sociocultural and political environments
- 2. Member of one or more professional nursing organizations and active at the state or national level in at least one professional nursing organization
- 3. Works with local, state and national legislators to influence the health care environment
- 4. Yearly continuing education
- 5. Certification in specialty area

Professional Maintenance L. Member of a professional nursing organization L. Yearly continuing education	3. 4.	Yearly continuing education Reads reviews of current research and evaluates their applicability to his/her own practice Participates in the research process by serving as a data collector, or subject in nursing research studies Mentoring of students and new nurses	<ol> <li>6.</li> </ol>	Reads and evaluates current nursing research for applicability into his/her own practice; puts relevant research findings into practice; works in conjunction with doctorally prepared nurses to design and implement clinically focused nursing research studies  Mentoring of students, RN's and new APN's	7.	Actively involved in outcome based clinical research and/or basic research to provide support for new nursing interventions, and to document the effectiveness of current nursing interventions Publishes and presents research and/or professional papers to expand the arena of nursing knowledge Mentoring of students, RN's, and various APNs

The expanding scope of knowledge and practice from different levels of nursing education and experience is demonstrated in figure 1. The square in the center represents the core values and basic knowledge which is shared by nurses of all educational backgrounds. The additional knowledge gained by the B.S.N. is demonstrated by a larger, all inclusive box. However, by the time the nurse reaches the master's level, clinical specialization begins to occur, represented by wedge shaped partitions out of the general area of master's level practice. The doctoral level is represented by an increasingly narrow focus, the pinnacle of which is the development of new knowledge. This also helps explain why nurses with doctorates (or doctoral students) frequently do not seem to be speaking the same language as anyone else, as each one focuses on an area in which he or she becomes an expert, to the temporary exclusion of other areas of nursing interest.

Figure 1



This continued expansion of nursing knowledge is assisted by theory and research, from nursing and other disciplines. As to use of theory, nursing as a profession needs theory to guide our own continuing development. Because much of what we do involves both the mind and the body, a mind-body framework for nursing research and care seems to be a logical choice. The following section describes PNI theory, on which this nursing model is based.

# Theoretical Framework - Psychoneuroimmunology and Therapeutic Nursing

Interventions: Of all the professionals who are working in the health care arena today, nurses are still the ones with the greatest amount of patient contact. Because of this, nurses are frequently in the best position to assess patients in a holistic manner, to see beyond just signs and symptoms and gain insight into the patient's mental and spiritual needs, as well as his/her physical needs. Nursing and nurses have long held the view that there is more to illness, and healing from illness, than can be explained by strictly medical/physiological theories of illness development and healing. These biological theories often leave many questions unanswered. Why is it that several people can be exposed to the same pathogen, but not everyone becomes sick in the same way? There are individual differences in patterns of disease development, but what are the factors that influence these patterns? Can stress and negative thoughts or feelings actually contribute to physical illness? And, can therapeutic nursing interventions, such as relaxation, support groups, and use of humor, help patients to become well, in addition to helping them feel well? Multi-disciplinary research in the area of psychoneuroimmunology (PNI) is just beginning to shed some new light on these old questions.

Psychoneuroimmunology started from a multifactorial model of illness which included stress, coping and disease formation (Engel, 1962). This theory was further developed by Solomon (1964, 1985, 1987) to include the effects of stress on the immune system in disease formation. Later, the term "psychoneuroimmunology" was coined by Ader (1981) to describe the basic phenomena of this theory - interactions between the nervous system and the immune system and the subsequent effects upon disease development and progression.

Because PNI theory acknowledges the multi-factorial nature of wellness and illness, it is particularly useful as a guide for nursing research and practice (Birney, 1991). PNI research is inherently multi disciplinary in scope, involving many of the health care professions.

So how can you as a nurse use PNI theory as a guide for your practice or research? Figure 2 is designed to be a model to base therapeutic nursing interventions on. The primary premise behind this model is that there are many factors which affect well-being, and not all of these factors are physical or genetic. Multiple factors such as stressful life events, personality factors, and behavioral factors, all have an effect upon the person's well-being. Whether this is a negative effect or a positive effect depends upon the event or the behavior. Psychosocial factors affect physical functioning through changes in the PNI network. The PNI network is an interconnection of reactions, involving neurotransmitters, neuropeptides, hormones, and immune system components (Pert, Ruff, Weber, and Herkendam, 1985; Pert, 1986). Studies supporting PNI theory and the interactive communication network between the CNS and the immune system have been reviewed in the nursing literature (Birney, 1991; Houldin, Lev, Prystowsky, Redei, and Lowery, 1991; Nguyen, 1991).

The second basic premise of this model is that psychosocial factors, such as negative life events, and physical factors, such as high blood sugar, can be effectively modified using the appropriate therapeutic nursing interventions. The first set of nursing interventions listed are the Complementary Nursing Interventions.

Complementary therapy involves incorporating various non-medical treatments into the total care of the patient.

Complementary interventions are not alternative therapies intended to replace traditional medical care, instead, they are used as additional or adjunctive treatments to improve patient mental and physical well-being. However, as noted by Lerner (1992), "any therapy that can be used adjunctively can and has been used alternatively, and vice versa", so there tends to be some overlap in the use of terms (p. 32). Complementary interventions, such as use of humor, relaxation therapy, imagery, hypnosis, massage, act to modify the effect of negative stressful life events, psychosocial factors, and behavioral factors upon the PNI network, and thus upon well-being.

#### **Psychosocial Factors** Stressful life events: Work related; Family related; **Physiological** Socio-Political Factors **Environmental Factors** role strain: etc. **Factors** Socio-Economic Status **Personality Factors Pathogens** Health Insurance Carcinogens Illness Cost of Health Care Depressive personality; coping Extremes of Heat, Cold, Humidity Injury Availability of health resources styles; Helplessness; **Environmental safety regulations Public Health conditions** Disabilities Hopelessness; Emotional Safe water, air and non-toxic environment Suppression etc. Physical safety from violence Social Support Family, Friends, Coworkers **Modifying Factors** Values **Traditional Interventions Modifying Factors Cultural values** Medical Care **Spiritual Values Nursing Care** Allied health interventions such as Political and social activism physical therapy, chiropractic etc. Increasing primary care providers **Modifying Factors** Efficient use of health resources Social support systems Health care reform **Public Health Modifying Factors** Sanitation **Immunizations Holistic Interventions** Regulatory Activism Mental health interventions Caring presence Communication Holistic Well-Being: Optimal physical; **Factors Not Subject to** PNI Network Modification mental; emotional Psycho-Neurological Functioning and spiritual Genetics **Neuro-Endocrine Functioning** Age functioning **Endo-Immunological Functioning Past Experiences**

The next set of interventions listed are the Environmental Interventions. These endeavors, such as professional activism for quality of care and safe staffing levels, or political activism to improve access to health care, are aimed at improving the health care environment. Improving the health care environment may not directly affect any one persons physical status, but act to improve the general environment and well-being of a person or a community of persons. In addition, environmental factors also can determine whether or not the person receives any intervention, from either complementary practitioners or traditional practitioners. Environmental interventions are most likely to be implemented by nurses with advanced educational levels, as they are usually more prepared to deal with these broad based environmental factors.

The third general set of interventions are the Traditional Interventions. These are primarily aimed at directly improving one person's physical functioning. Of course, physiological factors play an integral part in holistic well-being. In addition, physiological factors are those most of us are most comfortable in dealing with, and the ones most laypersons connect with health care. This may be because most traditional nursing, medical, and allied health interventions have been aimed at modifying these physiological factors directly.

The final set of factors I chose to include are the factors not subject to modification. There are only a few of these, but in some persons they profoundly affect well-being, so I felt the need to include these factors. However, while a person's genotype and past are not modifiable, how these things affect their physical and mental well-being may well be modifiable by any of the three general sets of interventions in this model. See table 2 for a beginning list of various therapeutic nursing interventions for each area.

Lastly, I would like to admit that many interventions may fall into more than one category. For example: individualized pain management, in my mind, is primarily a complementary intervention. Pain usually is considered a stressful life event, and can affect various psychosocial factors, behavioral factors and the PNI network. However, some types of pain directly affect physiological functioning, so you may consider pain management as a traditional intervention. There is certainly room for flexibility and growth in this model. Feel free to experiment with it to find what best fits your practice or research needs.

Table 2

Psychosocial Factors	Complementary Therapeutic Nursing Interventions	Environmental Factors	Environmental Therapeutic Nursing Interventions	Physiological Factors	Traditional Therapeutic Nursing Interventions
Stressful Life Events- Bereavement; Work-related; Family related; Role -strain; Personal Illness or Injury Personality Factors Personality; Self-Esteem, Depression; Fatigue; Coping Style; Helpless/ Hopelessness; Emotional Suppression; Social Support/Relationship Quality and quantity of social support from spouse, family, friends, and community. Quality of intimate relationships Religious/Cultural Factors: Religious beliefs/ cultural beliefs about health and health care, role of women, men, children in culture/society, behaviors/habits concerning nutrition and exercise, beliefs concerning use of health care system, beliefs about the appropriateness/effective ness of various medical and nursing interventions. Behavioral Factors: Smoking; Alcohol/Drug Use; Exercise; Nutrition; Sleep; Sexual Behavior;	Interventions which modify negative effects of various psychosocial factors upon mental health/PNI network/physical health, and thus improve well-being.  Nursing Interventions  Relaxation therapy Imagery Humor Massage Exercise Diet Support groups Hypnosis Touch Therapeutic Touch Aroma therapy Sound/Music therapy Yoga/Tai-Chi or other exercise/relaxation therapy Structured psychological interventions Family therapy Marriage therapy Crisis Intervention Spiritual retreats Other non-harmful alternative therapies that the client believes are helpful	Pathogens; Carcinogens; Socio-Economic Status; Cost of care; Funding Structures; Level of Unit Staffing; Working/Living Conditions; Public Health Conditions; Sanitation; Regulatory Controls; Access to Health Care; ect.	Those endeavors, such as professional activism for quality of care and safe staffing levels, or political activism to improve access to health care, which are aimed at improving the health care environment.  Nursing Interventions  Re-designing hospital and health care environments to meet client needs for a calm, healing environment  Developing critical pathways which improve client outcome and nursing effectiveness  Self-empowerment  Political and administrative activism  Lobbying  Belonging to a nursing organization which has a political action group  Networking with local and state lawmakers and business leaders to improve health care  Working within the system - Insurance companies, HMO's PPO's, to improve health care	Neurological; Cardiac; Skin; Digestive; Elimination; Reproductive; Respiratory; Regulatory; Musculoskeletal; Hematologic; Immune; Endocrine.	Primarily aimed at directly improving one person's physical functioning.  Nursing Interventions  Patient Education Physical support of clients, such as TCDB, early ambulation, ROM, etc. Basic Hygiene support Patient monitoring and documentation Nursing technical interventions such as PIC lines to prevent peripheral vein dysfunction; Appropriate use of IV pumps to avoid fluid overload; Cardiac monitoring to prevent injury related to cardiac arrthymias, etc. Appropriately administering routine and PRN medications Patient advocacy to obtain needed medical orders Various medical procedures under the direction of MD or APN

#### **Test of the Model**

Because this model is a midrange theory that covers most includes most types of nursing interventions, it can be used to guide nursing research in a number of areas. For instance, it could be used to test the effectiveness of either traditional or complementary nursing interventions at improving the healing rate in decubitus ulcers. It could also be used to determine the effectiveness of a community education program in reducing the number of cases of multiple drug resistant TB. Because of my own and local cancer patient's interest in use of humor, a small pilot study to determine the effectiveness of humor as a complementary nursing intervention is briefly described below.

#### **Purpose**

The purpose of this research is to determine the effect of mirthful laughter and sense of humor on stress and NK cytotoxicity in a sample of healthy adult females.

# **Significance**

The use of humor to decrease stress, pain, and perhaps influence immune functioning has recently become a popular complementary intervention (Gilligan, 1993; Groves, 1991; Simon, 1990; Sullivan & Deane, 1988). While relaxation therapy requires time to learn and practice, laughter in response to a humorous stimulus is a natural occurrence (Cogan, Cogan, Walts & McCue, 1987). In addition, use of humor does not take large amounts of time or money to implement. However, the effectiveness of humor in the reduction of stress, and the physiological effects of laughter, are just beginning to be supported in the literature. The effect of sense of humor and laughter on psychological and physiological functioning need further examination before widespread implementation as a complementary nursing intervention.

# **Research Question:**

Does Sense of Humor and/or exposure to a humor effect stress, anger, or Natural Killer (NK) cell activity? Theoretical Framework:

The Multifactorial Model for Holistic Nursing Intervention (Bennett 1994) was used to organize the concepts of humor as a holistic intervention, stress, anger and sense of humor as psychosocial factors, and NK cytotoxicity as a measure of immunological functioning.

# Methodology:

A pre-post test design was used. Sense of humor was measured using Martin and Lefcourt's (1984) Situational Humor Response Questionnaire (SHRQ), Coping Humor Scale (CHS) (Martin and Lefcourt, 1983) and the Multidimensional Sense of Humor Scale (MDHS) (Thorson and Powell, 1993). The State Anger Scale was used to measure anger (Spielberger 1983). The Stress-Arousal Check List (SACL) was used to determine perceived stress levels (Mackay, 1978). All of the psychological measures have acceptable reliability and validity using a college student sample. NK activity was measured using the standard chromium release assay.

# Hypotheses:

- 1. Exposure to a humorous video will enhance NK cell activity over baseline value.
- Sense of Humor will be correlated with NK baseline values, With subjects having higher humor scores also having higher baseline NK values.
- Sense of Humor will be correlated with increase in NK values at post movie and recovery time, with subjects having higher humor scores demonstrating greater increases in NK value following the humorous video.
- 4. Self-reported stress will decrease following the humorous video.
- 5. Self-reported anger will decrease following the humorous video.

The data were analyzed using Paired t-tests and Pearson's r.

# Sample:


Six healthy, adult,, female volunteers were solicited from the university setting for the first phase of this study.

# **Summary of Findings:**

There was no significant difference between pre NK and post NK levels (t = 0.78 p = 0.478). Scores on the humor scales did not significantly correlate with baseline NK values or with change in NK values. However, stress scores decreased significantly (t = 3.73 p = 0.014). In addition, anger decreased following the film (t = 2.39 p = 0.062). There was also an interesting correlation noted between reports of decreased stress (SACL) and NK activity following the intervention. Those subjects who reported the most decrease in stress following

the film, also had the most increase (or the least amount of decrease) in NK activity (r = .79 p = 0.111). This pilot study finding supports the premise that humor may affect the immune system through cognitive pathways, by reducing or modifying the effects of stress.

Figure 3



# **Summary:**

You may be wondering, "why go to all this effort to investigate humor? Humor is currently being used in a number of health care settings already." Nursing must not continue to develop patient care techniques by the process of trial and error, or simply use interventions because you just attended a workshop on that intervention. The role of the doctorally prepared nurse is to investigate and document the <u>effectiveness</u> of nursing interventions. Once we have adequate documentation of an interventions cost, scope of use, effectiveness and usual outcomes, it is much easier to acquire the funding needed to further implement that intervention in the appropriate cases. The pilot study as described above tested the effectiveness of a nursing intervention, humorous stimulus, at reducing stress and anger, and at improving immune system functioning. Admittedly, it was a very small pilot study, and additional research in this area is being planned. Hopefully, the

next study will support the model linkages found in the pilot study, and additional linkages may be determined.

Of course, these studies as outlined test only a small part of the model, so the results of the study can only be viewed in terms of those factors. However, this is the beginning of a series of studies to determine the effectiveness of various nursing interventions in improving immune functioning and/or well-being, in order to further test this model. See table 4 for a structure-process-outcome example of how this theory could be used and evaluated in a patient care setting.

In conclusion, this paper has described a model for nursing practice, and a plan for testing part of the model. In addition, a plan for estimating demand for services and funding to continue the research is proposed. Many of these plans are tentative, as this is the beginning phase of my own research career, in addition to being the dawn of a new model for nursing practice. Hopefully, the model and the plan will continue to develop throughout the coming years, into a truly useful model, and a successful research career.

# Table 4

Table 4							
Structure	Process	Outcome					
Institutional philosophy and policy acknowledges the effect of various psychosocial factors (life events, coping mechanisms, social support, personality traits) upon the well-being of all persons.	<ol> <li>Holistic nursing assessment is performed by the nurse to ascertain client physical, mental and spiritual needs.</li> <li>The client is informed of various treatment</li> </ol>	<ol> <li>The client verbalizes understanding of the effect of specific psychosocial factors on their well-being.</li> <li>Individual client needs are documented and appropriate plan of treatment is implemented.</li> </ol>					
The client is viewed as having an integral role the his/her healing process.	options and possible consequences, whenever possible.	Clients participate in the decision making process concerning their own plan of care.					
Institutional philosophy exists that supports use of complementary nursing interventions, in addition to standard therapeutic nursing interventions.	Appropriate therapeutic nursing interventions are implemented in accordance with client physical, mental, and spiritual needs.	Clients participate in therapeutic and complementary nursing interventions to moderate the effects of stress, as indicated by their individual needs.					
The educational preparation of the nurse includes instruction in the use of complementary and therapeutic nursing interventions.	Nurses maintain both technical and behavioral skills by continuing education and appropriate certification.	5. Client stress, (Daily Hassles Questionnaire) anxiety, (State-Trait Anxiety Inventory - STAI) and pain (McGill Pain Questionnaire) will be moderated by appropriate					
Policies and procedures reflect adequate flexibility to meet individual patient needs, including physical, mental and spiritual needs.	5. Persons skilled in various behavioral techniques, research methodology and statistical analysis are available to the nurse as consultants.	nursing interventions, as indicated by the clients individual needs.  6. Client immune function (Natural Killer Cell Activity-NKCA,					
6. Adequate resources are allocated to support complementary and therapeutic nursing care, including adequate staffing, supplies, equipment, and appropriate environment for healing.	Clients are treated as individuals with unique needs and capabilities.	Mitogen response) and coping abilities (Ways of Coping) will be enhanced by appropriate nursing interventions, as indicated by the client's individual needs.					
Nurses have sufficient autonomy and expertise to assess, implement and evaluate therapeutic nursing interventions.	7. Clients are instructed and supported in the use of various behavioral techniques to decrease the effects of life stressors upon their well-being, as indicated by the client's particular needs.	7. Client overall well-being (measured by Brief Symptom Inventory; Quality of Life Index; Center for Epidemiologic Studies Depression Scale, CES-D; The Hospital Anxiety and Depression Scale; STAI; NKCA; and increased survival time for those with metastatic cancer), will be					
Institutional policy, procedures, and resources support nursing research.	Nursing research documents client outcomes and cost-effectiveness of various nursing	enhanced through the use of individualized, therapeutic nursing interventions.					
9. Nurses performing therapeutic nursing interventions are supported by salaries appropriate to their level of expertise and by yearly educational allowances for course work and/or workshop attendance.	interventions.						

# References

- Ader, R., & Cohen, N. (1981). Conditioned immunopharmacologic responses. In R. Ader (Ed.), Psychoneuroimmunology (pp. 6-38). New York: Academic Press.
- Barlozzari, T., Leonhardt, J., Wiltrout, R., Herberman, R., & Reynolds, C. (1985). Direct evidence for the role of LGL on the inhibition of experimental tumor metastases. Journal of Immunology, 134, 2783-2789.
- Bartrop, R., Lazarus, L., Luckhurst, E., Kiloh, L., & Penny, R. (1977). Depressed lymphocyte function after bereavement. <u>Lancet</u>, <u>1</u>, 834-836.
- Bellert, J. (1989). Humor; a therapeutic approach in oncology nursing. Cancer Nursing, 12(2), 65-70.
- Berk, L., Tan, S., Nehlsen Cannarella, S., Napier, B., Lewis, J., Lee, J., & Eby, W. (1988). Humor associated laughter decreases cortisol and increases spontaneous lymphocyte blastogenesis. <u>Clinical Research</u>, <u>36</u>, 435A.
- Berk, L., Tan, S., Fry, W., Napier, B., Lee, J., Hubbard, R., Lewis, J., & Eby, W. (1989). Neuroendocrine and stress hormone changes during mirthful laughter. <u>The American Journal of the Medical Sciences</u>, <u>298</u>, 391-396.
- Birney, M. (1991). Psychoneuroimmunology: A holistic framework for the study of stress and illness. <u>Holistic Nursing Practice</u>, <u>5</u>(4), 32-8.
- Buchanan, L. (1994). Therapeutic nursing intervention knowledge development and outcome measures for advanced practice. <u>Nursing and Health Care</u>, <u>15</u>(4), 190-195.
- Cogan, R., Cogan, D., Walts, W., & McCue, M. . (1987). Effects of laughter and relaxation on discomfort thresholds. <u>Journal of Behavioral Medicine</u>, <u>10(2)</u>, 139-144.
- Cousins, N. (1979). Anatomy of an illness as perceived by the patient. Toronto: Bantam.
- Davidhizar, R., & Bowen, M. (1992). The dynamics of laughter. <u>Archives of Psychiatric Nursing</u>, <u>6</u>(2), 132-137.
- Dillon, K., Minchoff, B., & Baker, K. (1985). Positive emotional states and enhancement of the immune system. <u>International Journal of Psychiatric Medicine</u>, <u>15</u>, 13-18.
- Dorian, B., Keystone, E., Garfinkel, P., & Brown, G. (1981). Immune mechanisms in acute psychological stress. <u>Psychosomatic Medicine</u>, <u>43</u>, 84 (Abstr.).

- Dorken, H. (1994). The financing and organization of universal healthcare: A proposal for the National Academies of Practice. In <u>Healthcare utilization and cost series Vol. 3</u>. South San Fransico: Foundation for Behavioral Health.
- Engel, G. (1962). Psychological development in health and disease. Philadelphia: Saunders.
- Erdman, L. (1991). Laughter therapy for patients with cancer. Oncology Nursing Forum, 18(8), 1359-1363.
- Friedman, J., Trucker, J., Tomlinson-Keasey, C., & Schwartz, J. (1993). Does childhood personality predict longevity? <u>Journal of Personality and Social Psychology</u>, <u>65</u>(1), 176-185.
- Gilligan, B. (1993). A positive coping strategy: Humor in the oncology setting. <u>Professional Nurse</u>, <u>8</u>(4), 231-233.
- Glaser, R., Kiecolt-Glaser, J., Speicher, C., & Holliday, J. (1985a). Stress, loneliness, and changes in herpesvirus latency. Journal of Behavioral Medicine, 8, 249-260.
- Glaser, R., Mehl, V., Penn, G., & Speicher, C. (1986). Stress-associated changes in plasma immunoglobulin levels. <u>International Journal of Psychosomatics</u>, 33(2), 41-42.
- Glaser, R., Kennedy, S., Lafuse, W., & Bonneau, R. (1990). Psychological stress-induced modulation of interleukin 2 receptor expression and interleukin 2 production in peripheral blood leukocytes. <u>Archives of General Psychiatry</u>, <u>47</u>, 707-712.
- Groves, D. (1991). "A merry heart doeth good like a medicine.." Holistic Nursing Practice, 5(4), 49-56.
- Halley, F. (1991). Self-regulation of the immune system through biobehavioral strategies. <u>Biofeedback and Self-Regulation</u>, <u>16</u>(1), 55-74.
- Hanna, N., & Fidler, I. (1980). Role of natural killer cells in destruction of circulating tumor emboli. <u>Journal of the National Cancer Institute</u>, <u>65</u>, 801-809.
- Hanna, N., & Barton, R. . (1981). Definitive evidence that natural killer (NK) cells inhibit experimental tumor metastasis *in vivo*. <u>Journal of Immunology</u>, <u>127</u>, 1754-1758.
- Hanna, N. . (1986). *In vivo* activities of NK cells against primary and metastatic tumors in experimental animals. In E. Lotzova & R. Herberman (Eds.), <u>Immunobiology of natural killer cells</u>. Boca Raton, Florida: CRC.
- Henry, B., & Moddy, L. (1985). Energize with laughter. Nursing Success Today, 2(1), 5-8.

- Herberman, R., & Ortaldo, J. (1981). Natural killer cells: Their role in defenses against disease. <u>Science</u>, <u>214</u>, 24-30.
- Herth, K. (1984). Laughter; A nursing treatment. American Journal of Nursing, 84, 991-992.
- Houldin, A., Lev, E., Prystowsky, M., Redei, E., & Lowery, B. . (1991). Psychoneuroimmunology: A review of literature. <u>Holistic Nursing Practice</u>, <u>5</u>(4), 10-21.
- Hudak, D., Dale, A., Hudak, M., & DeGood, D. (1991). Effects of humorous stimuli and sense of humor on discomfort. <u>Psychological Reports</u>, <u>69(3)</u>, 779-786.
- Irwin, J., & Livnat, S. (1987). Behavioral influences on the immune system: Stress and conditioning. 9th Annual Meeting of the Canadian College of Neuro-Psychopharmacology (1986, Vancouver, Canada).

  Progress in Neuro-Psychopharmacology and Biological Psychiatry, 11(2-3), 137-143.
- Jemmott, J., Borysenko, J., Borysenko, M., McClelland, D., Chapman, R., Meyer, D., & Benson, H. (1984).

  Academic stress, power motivation, and decrease in secretion rate of salivary secretory immunoglobulin

  A. <u>Lancet</u>, 1, 1400-1402.
- Jemmott, J., & Magloire, K. (1988). Academic stress, social support, and secretory immunoglobulin A. Journal of Personality and Social Psychology, 55, 803-810.
- Kasl, S., Evans, A., & Niederman, J. (1979). Psychosocial risk factors in the development of infectious mononucleosis. <u>Psychosomatic Medicine</u>, <u>41</u>, 445-466.
- Keast, D. (1981). Immune surveillance and cancer. In K. Bammer & B. Newberry (Eds.), <u>Stress and Cancer</u> (pp. 71-97). Toronto: C.J. Hogrete.
- Kiecolt-Glaser, J., Ricker, D., Geroge, J., Messick, G., Speicher, C., Garner, W., & Glaser, R. (1984). Urinary cortisol levels, cellular immunocompetency, and loneliness in psychiatric inpatients. <a href="Psychosomatic">Psychosomatic</a>
  <a href="Medicine">Medicine</a>, 46, 15-23.
- Kiecolt-Glaser, J., Garner, W., Speicher, C., Penn, G., Holliday, J., & Glaser, R. (1984). Psychosocial modifiers of immunocompetence in medical students. <u>Psychosomatic Medicine</u>, <u>46</u>, 7-14.
- Kiecolt-Glaser, J., Glaser, R., Strain, E., Stout, J., Tarr, K., Holliday, J., & Speicher, C. (1986). Modulation of cellular immunity in medical students. <u>Journal of Behavioral Medicine</u>, 9, 5-21.

- Kiecolt-Glaser, J., Fisher, L., Ogrocki, P., Stout, J., Speicher, C., & Glaser, R. (1987a). Marital quality, marital disruption and immune function. <u>Psychosomatic Medicine</u>, 46, 13-34.
- Kiecolt-Glaser, J., Kennedy, S., Malkoff, S., Fisher, L., Speicher, C., & Glaser, R. (1988). Marital discord and immunity in males. <u>Psychosomatic Medicine</u>, <u>50</u>, 213-229.
- Kiessling, R., Klein, E., & Wigzell, H. (1975). "Natural" killer cells in the mouse. I. Cytotoxic cells with specificity for mouse Maloney leukemia cells. Specificity and distribution according to genotype. <u>European Journal of Immunology</u>, 5, 112-117.
- Killeen, M. (1991). Clinical clowning: Humor in hospice care. <u>American Journal of Hospice and Palliative</u>

  <u>Care, 8(3), 23-27.</u>
- Labott, S., Ahleman, S., Wolever, M., & Martin, R. . (1990). The physiological and psychological effects of the expression and inhibition of emotion. <u>Behavioral Medicine</u>, <u>16</u>(4), 182-189.
- Leidy, K. (1992). Enjoyable learning experiences- an aid to retention? <u>Journal of Continuing Education in Nursing</u>, <u>23(5)</u>, 206-208.
- Levy, S., Herberman, R., Maluish, A., Schlein, B., & Lipppman, M. (1985). Prognostic risk assessment in primary breast cancer by behavioral and immunological parameters. <u>Health Psychology</u>, 4, 99-113.
- Long, P. (1987). Laugh and be well? Psychology Today, 21(10), 28-29.
- MacHovec, F. (1991). Humor in therapy. <u>Psychotherapy in Private Practice</u>, <u>9</u>(1), 25-33.
- Martin, R., & Lefcourt, H. (1983). Sense of humor as a moderator of the relation between stressors and mood.

  <u>Journal of Personality and Social Psychology</u>, 45, 1313-1324.
- Martin, R., & Dobbin, J. (1988). Sense of humor, hassles, and immunoglobulin A: Evidence for a stress-moderating effect of humor. <u>International Journal of Psychiatry in Medicine</u>, <u>18</u>(2), 93-105.
- Martinez, J. (1989). The role of empathic humor in counteracting burnout and promoting renewal: The development and implementation of a personal growth workshop. Massachusetts: University of Massachusetts.
- McClelland, D., Ross, G., & Patel, V. (1985a). The effect of an academic examination on salivary

- norepinephrine and immunoglobulin levels. <u>Journal of Human Stress</u>, <u>11</u>, 42-59.
- McClelland, D., Davidson, R., & Saron, C. (1985). Stressed power motivation, sympathetic activation, immune function, and illness. <u>Advances</u>, <u>2</u>(2), 42-52.
- Metcalf, C. (1987). Humor, life, and death. Oncology Nursing Forum, 14(4), 19-21.
- Morreall, J. (1991). Humor and work. <u>Humor International Journal of Humor Research</u>, 4(3-4), 359-373.
- Mott, T. (1994). <u>American Journal of Clinical Hypnosis</u>. 2200 E. Devon Avenue, Suite 291 Des Plains IL 60018: The American Society of Clinical Hypnosis.
- Mouton, C., Fillion, L., Tawadros, E., & Tessier, R. (1989). Salivary IgA is a weak stress marker. <u>Behavioral Medicine</u>, <u>15</u>(4), 179-185.
- Nguyen, T. (1991). Mind, brain, and immunity: A critical review. Holistic Nursing Practice, 5(4), 1-9.
- Nieburg, H., Weiss, J., Navarrete, M., Strax, P., Teirstein, A., Grillione, C., & Siedlecki, B. (1979). The role of stress in human and experimental oncogenesis. Cancer Detection and Prevention, 2, 307-336.
- O'Leary, A. (1990). Stress, emotion, and human immune function. <u>Psychological Bulletin</u>, 108(3), 363-382.
- Osterlund, J. (1983). Humor A serious approach to patient care. Nursing, 12, 46-47.
- Palmblad, J., Cantell, K., Strander, H., Froberg, J., Karlsson, C., Levi, L., Granstom, M., & Unger, P. (1976). Stressor exposure and immunological response in man: Interferon-producing capacity and phagocytosis.

  <u>Journal of Psychosomatic Research</u>, 20, 193-199.
- Palmblad, J., Petrini, B., Wasserman, J., & Akerstedt, T. (1979). Lymphocyte and granulocyte reactions during sleep deprivation. <u>Psychosomatic Medicine</u>, <u>41</u>, 273-278.
- Parish, A. (1994). It only hurts when I don't laugh. American Journal of Nursing, 94(8), 46-48.
- Pert, C., Ruff, M., Weber, R., & Herkendam, M. (1985). Neuropeptides and their receptors: A psychosomatic network. <u>Journal of Immunology</u>, <u>135</u>, 820-826.
- Pert, C. (1986). The wisdom of the receptors: Neuropeptides, the emotions, and body mind. <u>Advances</u>, <u>3</u>(3), 8-16. Raber, W. (1987). The caring role of the nurse in the application of humor therapy to the patient experiencing helplessness. <u>Clinical Gerontologist</u>, <u>7</u>(1), 3-11.
- Robinson, V. (1979). Humor in nursing. In C. Carlson & B. Blackwell (Eds.), <u>Behavioral Concepts and Nursing Intervention</u>. Philadelphia: Lippincott.

- Ruxton, J., & Hester, M. (1987). Humor: Assessment and interventions. Clinical Gerontologist, 7(1), 13-21.
- Ruxton, J. (1988). Humor intervention deserves our attention. Holistic Nursing Practice, 2(3), 54-62.
- Samra, C. (1985). A time to laugh. <u>Journal of Christian Nursing</u>, <u>3</u>, 15-19.
- Saper, B. (1988). Humor in psychiatric healing. Psychiatric Quarterly, 59(4), 306-319.
- Schleifer, S., Keller, S., Camerion, M., Thornton, J., & Stein, M. (1983). Suppression of lymphocyte stimulation following bereavement. JAMA, 250, 374-377.
- Schmidt, R., Bartley, G., Levine, H., Schlossmann, S., & Ritz, J. (1985). Functional characterization of LFA-1 antigens in the interaction of human NK clones and target cells. <u>Journal of Immunology</u>, <u>135</u>, 1020-1025.
- Scott, C. (1985). Go ahead, laugh, it's good for you. Nursing Success Today, 2(1), 8.
- Simon, J. (1988). Therapeutic humor: Who's fooling who? <u>Journal of Psychosocial Nursing and Mental</u>

  <u>Health Services</u>, <u>26</u>(4), 8-12.
- Simon, J. (1989). Humor techniques for oncology nurses. Oncology Nursing Forum, 16(5), 667-670.
- Simon, J. (1990). Humor and its relationship to perceived health, life satisfaction, and morale in older adults.

  <u>Issues in Mental Health Nursing</u>, 11(1), 17-31.
- Smith, K. (1992). Optimism, self-consciousness, and coping with aids and cancer. California: Pacific Graduate School of Psychology.
- Solomon, G., & Moos, R. (1964). Emotions, immunity, and disease: A speculative theoretical integration.

  Archives of General Psychiatry, 11, 657-674.
- Solomon, G. (1985). The emerging field of psychoneuroimmunology. Advances, 2, 6-19.
- Solomon, G. (1987a). Psychoneuroimmunology: Interactions between central nervous system and immune system. <u>Journal of Neuroscience Research</u>, <u>18</u>, 1-9.
- Steinhauer, E., Doyle, A., Reed, J., & Kadish, A. (1982). Defective natural cytotoxicity in patients with cancer:

  Normal number of effector cells but decreased recycling capacity in patients with advanced disease.

  <u>Journal of Immunology</u>, 129, 2255-2259.
- Stone, A., Cox, D., Valdimarsdottir, H., & Neale, J. (1987). Secretory IgA as a measure of immunocompetence. Journal of Human Stress, 13(3), 136-140.
- Sullivan, J., & Deane, D. (1988). Humor and health. Journal of Gerontological Nursing, 14(1), 20-24.

- Thorson, J., & Powell, F. (1993). Development and validation of a multidimensional sense of humor scale.

  <u>Journal of Clinical Psychology</u>, 49(1), 13-23.
- Trent, B. (1990). Ottawa lodges add humor to armamentarium in fight against cancer. <u>Canadian Medical Association Journal</u>, <u>142(2)</u>, 163-166.
- U.S. Department of Health and Human Service. (1987). <u>Surgeon General's report on acquired immune</u>

  <u>deficiency</u> <u>syndrome</u>. Washington, DC: Author.
- Williams, H. (1986). Humor and healing: Therapeutic effects in geriatrics. Gerontion, 1(3), 14-17.