



ABSTRACT FORM

This form should be read in conjunction with the **ISEI ABSTRACT GUIDELINES**.

These 2 pages have been provided separately to assist you with the submission of your Abstracts in Word format to the Conference Organising Committee and online to the *International Journal of Exercise Science* (IJES).

You **MUST** complete this entire Abstract Form to provide us with details needed to assign your abstract to the correct theme.

Also advise us if you are applying to be considered for one of the Early Career Researcher Awards (Best Poster and Best Oral Presentation).

Send this entire form by E-mail to the ISEI Organising Committee, but **ALSO** follow the guidelines for submission of the Abstract (single page only) to the *IJES*.

ABSTRACT SUBMISSION – DEADLINE 10 May 2013

Title (up to 30 words, Arial, 11 pt, single line spaced, in sentence case. Like this:	Exhaustive exercise enhances immune response to flagellin via adrenaline-mediated up-regulation of TLR5 expression
Authors (Underline the <u>presenting author</u>)	<u>M Uchida</u> , E Oyanagi, M Miyachi, MJ Kremenik, S Onodera & H Yano
Department, Institution, Country	Division of Health Science, Kawasaki University of Medical Welfare, Okayama, Japan
Address Corresponding Author	288 Matsushima, Kurashiki, OK 701-0193, Japan, and E-mail adress: masa19850525@hotmail.co.jp
Select Your Abstract Session Theme Category	5 poster session
Preferred Presentation Form	<input type="checkbox"/> Oral communication <input checked="" type="checkbox"/> Poster Note final decisions on format of presentation will be by the ISEI Scientific Committee
Is the presenter eligible for the Early Career Researcher Awards (poster and oral awards)? (Previous winners are ineligible for same category)	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No Eligible persons are those studying for a higher degree – MSc, MPhil or PhD – or who have completed their PhD within the last 3 years as at 10th September 2013



***ISEI Abstract – Please do not change any fonts, font sizing, or spacing!
Replace this wording with your selected “Session theme number – 5”***

Exhaustive exercise enhances immune response to flagellin via adrenaline-mediated up-regulation of TLR5 expression

UCHIDA M1, OYANAGI E2, MIYACHI M2, KREMENIC MJ3, ONODERA S3, and YANO H3.

1Division of Health Science, and 3Department of Health and Sports Science, Kawasaki University of Medical Welfare, Okayama, Japan, and 2Department of Health Promotion and Exercise, National Institute of Health and Nutrition, Tokyo, Japan

ABSTRACT

Objective: We already reported that lipopolysacchride (LPS)-induced tumor necrosis factor (TNF)- α production as a bacterial infection model to induce immune response, was inhibited by exhaustive exercise. However, it remains unclear whether or not the immune response to flagellin (FG), which binds to toll like receptor 5 (TLR5) and induces pro-inflammatory cytokine production, is also inhibited by this severe exercise. The aim of this study was to determine whether or not exhaustive exercise affects TNF- α productions after FG injection in mice. **Methods:** Both exhaustive-exercised (EX; n=12) and non-exercised (N-EX; n=12) male C3H/HeN mice were injected with FG (1 mg/kg, i.v), and blood samples were collected. In addition, to clarify the effect of catecholamine on immune response macrophage and intestinal cells after FG stimulation, RAW264 cells and Caco2 cells were cultured 30min after propranolol (Prop; β -adrenergic receptors blocker) or Ly294002 (Ly; PI3K inhibitor) treatments, and were then stimulated with adrenaline (AD; 1 μ M) and FG (5 μ g/ml). Moreover, the effect of Prop (10 mg/kg, n=12) on FG-induced TNF- α production in EX mice was also examined. **Results:** TNF- α in EX group was significantly higher than that in N-EX group after FG injection. In epithelium cells, more intensity of TLR5 localization was observed on the plasma membrane area than in the cytosol area in EX mice, but not N-EX mice. Caco2 cells, but not RAW264 cells, significantly increased the FG-induced TNF- α production using AD treatment. Moreover, Prop treatment attenuated the AD-induced TNF- α production in response to FG in Caco2 cells. Although TLR5 expression on RAW264 cells was significantly decreased after AD treatment, the expression on Caco2 cells was rapidly increased. In fact, we observed the AD-dependent TLR5 translocation from cytoplasm to cell membrane in Caco2 cells, and the membrane translocation was inhibited by Prop and Ly treatment. Moreover, the pretreatment with Prop attenuated the exercise-induced plasma TNF- α response to FG in vivo. **Conclusion:** Our results suggest that immune response to FG via TLR5 might be enhanced by exhaustive exercise in mice.