

January 2008

Biogenic Porosity and its Lattice Boltzmann Method Permeability in the Karst Biscayne Aquifer

Kevin Cunningham
US Geological Survey

Mike Sukop
Florida International University


Haibo Huang
Florida International University

Pedro Alvarez
Florida International University

Allen Curran
Florida International University

See next page for additional authors

Follow this and additional works at: http://digitalcommons.wku.edu/geog_fac_pub

 Part of the [Fresh Water Studies Commons](#), [Geology Commons](#), and the [Geophysics and Seismology Commons](#)

Recommended Repository Citation

Cunningham, Kevin; Sukop, Mike; Huang, Haibo; Alvarez, Pedro; Curran, Allen; Waker, Michael; Florea, Lee J.; Renken, Robert; and Dixon, Joann F. (2008). Biogenic Porosity and its Lattice Boltzmann Method Permeability in the Karst Biscayne Aquifer. *Karst From Recent to Reservoirs*, 30-35.

Available at: http://digitalcommons.wku.edu/geog_fac_pub/4

Authors

Kevin Cunningham, Mike Sukop, Haibo Huang, Pedro Alvarez, Allen Curran, Michael Waker, Lee J. Florea, Robert Renken, and Joann F. Dixon

Biogenic Macroporosity and Its Lattice Boltzmann Method Permeability in the Karst Biscayne Aquifer

Kevin J. Cunningham

*U.S. Geological Survey
3110 SW 9th Avenue
Fort Lauderdale, FL 33315
United States of America
kcunning@usgs.gov*

Michael C. Sukop

Haibo Huang

Pedro F. Alvarez

*Department of Earth Sciences
Florida International University
11200 SW 8th Street
Miami, FL 33199
United States of America
sukopm@fiu.edu
huangh@fiu.edu
palva003@fiu.edu*

H. Allen Curran

*Department of Geology
Smith College
44 College Lane
Northampton, MA 01063
United States of America
acurran@email.smith.edu*

Michael A. Wacker

Lee J. Florea

Robert A. Renken

Joann F. Dixon

*U.S. Geological Survey,
3110 SW 9th Avenue
Fort Lauderdale, FL 33315
United States of America*

mwacker@usgs.gov, lflorea@usgs.gov, rarenken@usgs.gov, jdixon@usgs.gov

We focus on two major problems in the study of paleokarst of the Biscayne aquifer in southeastern Florida: (1) current conceptual models of karst aquifers do not adequately characterize much of the eogenetic macropore system within the carbonate rocks of the Biscayne aquifer, and (2) standard laboratory core-analysis methods cannot be used to accurately measure the permeability of highly macroporous carbonate core samples.

Cyclostratigraphic and borehole geophysical analyses of continuous coreholes (Cunningham et al., 2004; 2006a,b), trac-

er-test analyses (Renken et al., 2005), high-resolution X-ray computed tomography (HRXCT), and flow simulations using lattice Boltzmann methods (Sukop and Thorne, 2006) were used in combination to characterize the macroporosity and permeability of the eogenetic karst limestone of the Biscayne aquifer, principally within the 246-km² study area (Figure 1) of Cunningham, et al. (2004; 2006a,b). Results show that tabular-shaped, touching-vug hydrogeologic units are commonly stratiform within high-frequency cycles. Much of this macroporosity is biogenic in origin and concentrates groundwater flow to form the dominant type of flow passageway (Figure 2). Biogenic




Figure 1. Maps of study area. A. General location within the United States. B. Main study area of Cunningham et al. (2004; 2006a,b) in southeastern Florida and focus of this paper. C. Location of Northwest Wellfield within main study area. D. Location of cross-section A-A' (Figure 2.) within the Northwest Wellfield.

macroporosity largely manifests as two forms of touching-vug macroporosity: (1) ichnogenic macroporosity primarily related to post-depositional burrowing activity by callianassid shrimp (Figure 3); and (2) biomoldic macroporosity originating from the dissolution of fossil hard parts, principally mollusk shells. Ichnogenic macroporosity constrains the spatial distribution of at least some of the near-surface small caves (Figure 4) known to exist within the upper Pleistocene limestone of the Biscayne aquifer (Cressler, 1993). Biogenic macroporosity represents an alternative pathway for concentrated groundwater flow that differs considerably from the modern paradigm of karst aquifers that describe groundwater movement through cross-stratal dissolution features, bedding-plane vugs, and fractures (White, 1988; Ford and Williams, 1989; Worthington et al., 2000). Al-

though some of these more typical karst features occur in the study area, they less important than the biogenic flow zones.

The permeability of macroporous karstic rocks is extremely difficult to measure in a laboratory setting due to flow-rate limitations of the measuring apparatus, and issues related to maintaining and measuring the extremely small gradients needed to sustain Darcian flow regimes. HRXCT methods can produce digital reconstructions of porous media that are essential for detailed pore-scale flow modeling. The HRXCT provided rendering data for lattice Boltzmann calculation of permeability in samples with well-connected macropores.

Seven samples representative of biomoldic and trace-fossil




Figure 2. Cross-section A-A' showing vertical and lateral arrangement of biogenic macroporosity (fossil-moldic and ichnogenic macroporosity) and matrix porosity in the context of a high-frequency cyclostratigraphy for the Fort Thompson Formation and Miami Limestone of the Biscayne aquifer, Northwest Well Field (Figure 1.). Stratiform zones of substantial ichnogenic macroporosity interpreted to be related principally to callianassid burrowing are shown in light gray. Modified from Cunningham et al. (2006b).




Figure 3. Examples of ichnogenic macroporosity (intra- and inter-burrow) formed by the dense occurrence of *Ophiomorpha* shafts and tunnels within limestone of the Biscayne aquifer, Miami, Florida (Figure 1.). A. Photograph was acquired by a digital-optical borehole-image tool scanning the borehole wall of monitoring well G-3849 (Figure 1B.) in the Fort Thompson Formation (Middle Pleistocene, Marine Isotope Stage 11). The borehole wall image has been unwrapped to display an entire 360° view. Throughout the vertical interval, the borehole has a diameter of about 20.3 cm and an approximate circumference of 65 cm. Modified from Cunningham and Curran (2007). B. Ichnogenic macroporosity along cave wall inside Fat Sleeper Cave located about 26 km to the southeast of test corehole G-3849. The cave is developed in peloid-oid grainstone of the Late Pleistocene, Marine Isotope Stage 5e part of the Miami Limestone. The grainstone is overprinted with a maximum-intensity *Ophiomorpha* ichnofabric. The scale and distribution of cm-scale ichnogenic macroporosity on the walls of the cave are comparable to that shown on the borehole walls of test corehole G-3849. Photo by Allan M. Cressler.

related macroporosity were collected from the Fort Thompson Formation and Miami Limestone of the Biscayne aquifer in southeastern Florida. These samples were scanned using HRXCT methods that provided resolution on the order of 0.3 mm per pixel. Permeability calculations were conducted using lattice Boltzmann techniques applied to seven renderings of samples created from the HRXCT scans. Permeability was only calculated for the macroporous part of the seven samples and not the matrix porosity, because the rock matrix was simulated as a solid material in the computer renderings. Non-Darcian inertial flows were avoided by applying extremely small gradients while making the permeability calculations. The lattice Boltzmann method was verified against analytical solutions for

pipe and conduit flow. Measured horizontal permeabilities derived from lattice Boltzmann methods correspond to hydraulic conductivity values ranging from 0 m/s (for a sample lacking well-connected macroporosity) to 136 m/s, and are as much as five orders of magnitude greater than the largest values documented in laboratory tests.

REFERENCES

Cressler, A., 1993, The caves of Dade County, Florida: in *Georgia Underground: Dogwood City Grotto, Inc., of the National Speleological Society*, v. 30, no. 3, p. 9-16.




Figure 4. Inside Fat Sleeper Cave located near the shore of Biscayne Bay and about 17 km to the southeast of the main study area (Figure 1.) of Cunningham et al. (2004; 2006a,b). The cave is developed in peloid-oid grainstone overprinted with a maximum-intensity callianassid ichnofabric dominated by *Ophiomorpha*. White box inset is displayed in Figure 3B. The callianassid ichnofabric dominates a widely occurring stratiform layer within the Miami Limestone. Photo by Allan M. Cressler

Cunningham, K.J., and Curran, H.A., 2007, Ichnogenic porosity, high-frequency cyclostratigraphy, and groundwater flow in the karst Biscayne aquifer, SE Florida, USA, in Zonneveld, J.-J., and Gingras, M.K., eds., *International Ichnofabrics Workshop IX Abstracts with Program*, Calgary, Alberta, Canada, p. 12-15.

Cunningham, K.J., Carlson, J.L., Wingard, G.L., Robinson, Edward, and Wacker, M.A., 2004, Characterization of aquifer heterogeneity using cyclostratigraphy and geophysical methods in the upper part of the karstic Biscayne aquifer, southeastern Florida: U.S. Geological Survey Water-Resources Investigation Report 03-4208, 46 p.

Cunningham, K.J., Renken, R.A., Wacker, M.A., Zygnerski, M.R., Robinson, E., Shapiro, A.M., and Wingard, G.L., 2006a, Application of carbonate cyclostratigraphy and borehole geophysics to delineate porosity and preferential flow in the karst limestone of the Biscayne aquifer, SE Florida, in Harmon, R.S., and Wicks, C., eds., *Perspectives on karst geomorphology, hydrology, and geochemistry—A tribute volume to Derek C. Ford*

and William B. White: Geological Society of America Special Paper 404, p. 191-208, doi:10.1130/2006.2404(16).

Cunningham, K.J., Wacker, M.A., Robinson, Edward, Dixon, J.F., and Wingard, G.L., 2006b, A cyclostratigraphic and borehole geophysical approach to development of a three-dimensional conceptual hydrogeologic model of the karstic Biscayne aquifer, southeastern Florida: U.S. Geological Survey Scientific Investigations Report 2005-5235, 69 p.

Ford, D.C., and Williams P.W., 1989, *Karst geomorphology and hydrogeology*: London, Unwin Hyman, 601 p.

Renken, R.A., Shapiro, A.M., Cunningham, K.J., Harvey, R.W., Metge, D.W., Zygnerski, M.R., Osborn, C.L., Wacker, M.A., and Ryan, J.N., 2005, Assessing the vulnerability of a municipal well field to contamination in a karst aquifer: *Environmental and Engineering Geoscience*, v. 11, no. 4, p. 341-354.

Sukop, M.C., and Thorne, D.T., Jr., 2006, *Lattice Boltzmann modeling: An introduction for geoscientists and engineers*: New York, Springer, 172 p.

White, W.B. 1988, *Geomorphology and hydrology of karst terrains*: New York, Oxford University Press, 464 p.

Worthington, S.R.H., Ford, D.C., and Davies, G.J., 2000, Matrix, fracture and channel components of storage and flow in a Paleozoic limestone aquifer: in Sasowsky, I.D., and Wicks, C.M., eds., *Groundwater flow and contaminant transport in carbonate aquifers*, Rotterdam, Balkema, p. 113-128.

ecial

xon,
pore-
men-
ayne
cien-

ology

R.W.,
M.A.,
unici-
men-

mann
neers: