

Feb 14th, 2:50 PM

# Contribution of Freshwater Bivalves to Muskrat Diets in the Green River, Mammoth Cave National Park, Kentucky

Kimberly Asmus Hersey  
*University of Tennessee, Knoxville*

Joseph D. Clark  
*Southern Appalachian Research Branch, U.S. Geological Survey*

James B. Layzer  
*Tennessee Cooperative Fishery Research Unit, U.S. Geological Survey*

Follow this and additional works at: [http://digitalcommons.wku.edu/mc\\_reserch\\_symp](http://digitalcommons.wku.edu/mc_reserch_symp)

 Part of the [Animal Sciences Commons](#), [Forest Sciences Commons](#), [Geology Commons](#), [Hydrology Commons](#), [Other Earth Sciences Commons](#), and the [Plant Sciences Commons](#)

---

## Recommended Citation

Kimberly Asmus Hersey, Joseph D. Clark, and James B. Layzer, "Contribution of Freshwater Bivalves to Muskrat Diets in the Green River, Mammoth Cave National Park, Kentucky" (February 14, 2013). *Mammoth Cave Research Symposia*. Paper 15. [http://digitalcommons.wku.edu/mc\\_reserch\\_symp/10th\\_Research\\_Symposium\\_2013/Research\\_Posters/15](http://digitalcommons.wku.edu/mc_reserch_symp/10th_Research_Symposium_2013/Research_Posters/15)

This is brought to you for free and open access by TopSCHOLAR®. It has been accepted for inclusion in Mammoth Cave Research Symposia by an authorized administrator of TopSCHOLAR®. For more information, please contact [topscholar@wku.edu](mailto:topscholar@wku.edu).

# Contribution of Freshwater Bivalves to Muskrat Diets in the Green River, Mammoth Cave National Park, Kentucky

Kimberly Asmus Hersey<sup>1</sup>, Joseph D. Clark<sup>2</sup>, James B. Layzer<sup>3</sup>

<sup>1</sup> Department of Forestry, Wildlife and Fisheries, University of Tennessee, Knoxville

<sup>2</sup> Southern Appalachian Research Branch, U.S. Geological Survey

<sup>3</sup> Tennessee Cooperative Fishery Research Unit, U.S. Geological Survey

## Abstract

Musk rats (*Ondatra zibethicus*) are known to prey on freshwater bivalves (mussels and clams) and can negatively impact imperiled mussel species. However, factors that influence muskrat predation on bivalves are poorly understood. We evaluated the feeding ecology of muskrats at Mammoth Cave National Park, Kentucky by using stable isotope analysis of muskrat hair samples and by monitoring bivalve shell deposition at muskrat middens. Bayesian mixing-model analysis of stable isotope  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  ratios revealed that the median muskrat biomass derived from bivalves was 51.4% (5th and 95th percentiles were 39.1 to 63.4%, respectively), a much higher dietary proportion than previously reported. Shell depositions by muskrats at middens decreased with the availability of seasonal emergent vegetation, suggesting that the consumption of animal matter is in response to a scarcity of plant foods. Our results add to the growing body of evidence that muskrats have the potential to impact mussel population growth and recovery in some environments.