

Low and moderate intensity strength exercise affects more beneficially the lipid profile than high intensity strength exercise

#55

Fábio Santos Lira^{1,2}, Alex Shimura Yamashita¹, Marco Carlos Uchida^{1,3}, Nelo Eidy Zanchi⁴, Bruno Gualano⁴, Eivor Martins Jr¹, Marília Seelaender¹.

¹Molecular Biology of the Cell Group, Institute of Biomedical Sciences, Department of Cell and Developmental Biology, University of São Paulo, Brazil; ²Department of Physiology, Division of Nutrition Physiology, Federal University of Sao Paulo, Sao Paulo, Brazil; ³School of Physical Education - UNIFIEO, Osasco, Brazil; ⁴Laboratory of Applied Nutrition and Metabolism, Physical Education and Sport School, University of São Paulo, Brazil.

E-mail: fiolira@usp.br

This study aimed to compare the time-course effects of four different intensities of strength exercise (bench press) bouts on the blood lipid profile. Thirty-five Brazilian Army male soldiers were allocated randomly into five groups based at different percentages of one repetition maximum, in previous test (1-RM): 50%-1RM, 75%-1RM, 90%-1RM, 110%-1RM (this executing only eccentric phase), and control group. The total volume (sets x reps x load) of the exercise was equalized. The lipid profile (Triglycerides, VLDL, HDL-cholesterol, LDL-cholesterol, HDL-c/Total Cholesterol ratio and Total cholesterol) was determined at rest and after 1, 24, 48 and 72 h of the strength exercise. The 75% group demonstrated greater TG and VLDL reduction when compared with the other groups ($p<0.05$). Additionally, the 110% group presented an increased TG and VLDL concentration when compared with the control, 50% and 75% groups ($p<0.05$). HDL-c concentration was significantly greater after strength exercise at 50% and 75% when compared with 110% ($p<0.05$). Accordingly, the 50% and control groups had greater HDL-c concentration than 110% group after 48 h and 72 h ($p<0.05$). Finally, The 50% group showed lesser LDL-c concentration than 110% group after 24 h ($p<0.05$). No significant differences were found in Total Cholesterol and HDL-c/Total cholesterol ratio concentration. Results indicate that acute strength exercise changes lipid profile in a specific-intensity manner. Overall, low and moderate exercise intensities appear to promote more benefits on lipid profile than high intensity. Long term studies should confirm these findings.

Key words: **lipid profile; strength exercise; exercise intensity.**