

## VO<sub>2</sub> Reserve vs. Heart Rate Reserve During Moderate Intensity Treadmill Exercise

TANNER J. SOLHEIM\*, BRAD G. KELLER\*, and CHARLES J. FOUNTAINE‡

Department of Health, Physical Education and Recreation, University of Minnesota Duluth, Duluth, MN, USA

\*Denotes undergraduate student author, ‡Denotes professional author

---

### ABSTRACT

*International Journal of Exercise Science 7(4) : 311-317, 2014.* VO<sub>2</sub> and heart rate (HR) are widely used when determining appropriate training intensities for clinical, healthy, and athletic populations. It has been shown that if the % reserve (%R) is used, rather than % of max, HR and VO<sub>2</sub> can be used interchangeably to accurately prescribe exercise intensities. Thus, heart rate reserve (HRR) can be prescribed if VO<sub>2</sub> reserve (VO<sub>2</sub>R) is known. Therefore, the purpose of this study was to compare VO<sub>2</sub>R and HRR during moderate intensity exercise (50%R). Physically active college students performed a maximal treadmill test to exhaustion. During which VO<sub>2</sub> and HR were monitored to determine max values. Upon completion of the maximal test, calculations were made to determine the % grade expected to yield approximately 50% of the subjects VO<sub>2</sub>R. Subjects then returned to complete the submaximal test (50%R) at least two days later. The %VO<sub>2</sub>R and %HRR were calculated and compared to the predicted value as well as to each other. Statistical analysis revealed that VO<sub>2</sub> at 50%R was significantly greater than the actual VO<sub>2</sub> achieved,  $p < .001$ . Conversely, the mean predicted HR at 50%R was significantly less than the actual HR achieved,  $p < .001$ . In conclusion, this study indicated that VO<sub>2</sub> could be more accurately predicted than HR during moderate intensity exercise. The weak correlation between VO<sub>2</sub>R and HRR indicates that caution should be used when relying on a HR to determine VO<sub>2</sub>.

KEY WORDS: Exercise prescription, college students, submaximal testing

### INTRODUCTION

Heart rate (HR) and VO<sub>2</sub> are widely used to assess aerobic capacities and prescribe exercise programs to clinical, healthy, and athletic populations. However, it is important to be able to differentiate between % maximal oxygen uptake (%VO<sub>2max</sub>), % oxygen uptake reserve (%VO<sub>2</sub>R), % heart rate max (%HR<sub>max</sub>), and % heart rate reserve (%HRR). This is due to the fact that VO<sub>2</sub> and HR do not have absolute zeros, and their max values vary

depending on the individual, thus %VO<sub>2max</sub> and %HR<sub>max</sub> cannot be directly correlated. However, the term reserve describes the difference between the maximum and resting value for a given measurement providing more accurate predictive capabilities. Research has supported the finding that %VO<sub>2</sub>R and %HRR are more closely correlated than %VO<sub>2max</sub> and %HR<sub>max</sub> (5). This association enables more precise exercise prescription.

This association between %VO<sub>2</sub>R and %HRR was first demonstrated in 1997 with the use of cycle ergometers and it was concluded that %VO<sub>2</sub>R and %HRR should be used when prescribing exercise intensities. (7). The same relationship was found a year later with the use of treadmills. When exercise intensities were prescribed using a percentage of VO<sub>2</sub>R, the same percentage of HRR was observed (8). Further research in this field has provided evidence that indicates this association to be present on a much larger scope. For example, this association was found to be present in elliptical exercise (4), as well as in individuals with substandard health, such as those suffering from heart disease (1). All of these studies looked at maximal tests, from which, regression lines were constructed to determine the association between %VO<sub>2</sub>R and %HRR. These experiments made the assumption that this regression held true throughout all intensity levels. However, the majority of research has not been conducted at moderate intensities to accurately determine the relationship between %VO<sub>2</sub>R and %HRR at these levels.

Currently exercise programs are prescribed using either %VO<sub>2</sub>R or %HRR and the American College of Sports Medicine (ACSM) provides guidelines of specific percentages for varying intensities (9). If earlier findings are correct then a subject exercising at a set %VO<sub>2</sub>R should exhibit the same %HRR, and vice-versa. However, recent research has been contradictory to earlier findings that a 1:1 ratio exists at all exercise intensities. It has been found that as exercise intensity increases, %VO<sub>2</sub>R and %HRR deviate from the expected 1:1 ratio (3). It is plausible to assume that this

phenomenon exists during moderate exercise. Therefore, the purpose of this study was to determine if a 1:1 relationship exists between %HRR and %VO<sub>2</sub>R during moderate (50%) treadmill exercise.

## METHODS

### *Participants*

28 college students (11 male, 17 female, age 21.9 ± 1.6 years) participated in this study. The University's Institutional Review Board approved this study and all participants were presented with an informed consent form as well as a Physical Activity Readiness Questionnaire (PAR-Q) prior to testing. If potential participants answered 'yes' to any of the questions on the PAR-Q, this resulted in automatic disqualification from the study. The informed consent highlighted the purpose of the study and what was expected of the participants.

### *Protocol*

Subjects were required to perform two separate exercise bouts on two different days with at least 48 hours of rest between successive bouts. Weight (kg) was taken prior to each exercise bout using a Befour portable scale (Befour, Inc. Saukville, Wisconsin).

The first exercise bout was a treadmill VO<sub>2</sub> max test to exhaustion. The protocol was the "12 kmh/7.5 mph VO<sub>2</sub>max Test" found in the COSMED Fitmate user manual, NH Edition (COSMED Srl - Italy) (Table 1). The procedure was programmed into a Quinton Q-Stress™ Stress Test System that was synced with a TM55 Treadmill (Quinton Instruments, Bothell, WA).

In order to collect a minute by minute recording of HR, a Polar heart rate monitor (Polar Electro, Tempere, Finland) was lubricated using Spectra® 360 Electrode Gel (Scientific Digital Imaging, LLC, Oconomowoc, WI) and fitted to the subject. Minute-by-minute recordings of VO<sub>2</sub> were collected via indirect calorimetry with the use of a metabolic cart (Parvo Medics True One 2400, Sandy UT, USA). To ensure complete analysis of expired gases a nosepiece was worn in addition to a mouthpiece that was held in place using a light headpiece.

The protocol was explained before the start of the test (Table 1). The subjects were encouraged to perform until absolute exhaustion at which point the participants were instructed to straddle the belt of the treadmill while it was brought to rest. At the completion of the test, VO<sub>2max</sub> and HR<sub>max</sub> were collected from the PARVO system's minute-by-minute printout and imported into Microsoft® Excel® for future analysis. An RER of > 1.1 was considered to indicate that a true max was achieved.

On a separate day, at least 48 hours following the initial test, subjects returned to the lab to complete an exercise bout on a treadmill at an intensity that was predicted to yield 50% of the participants VO<sub>2</sub> reserve. This was calculated using the following equation:  $50\%VO_2 = (VO_{2max} - 3.5\text{mL/kg/min}) \cdot 50\% + 3.5\text{mL/kg/min}$ . Once 50%VO<sub>2</sub> was calculated, exercise intensity was determined using a speed of 3 mph (80.4m/min) with varying degrees of incline, using the following equation:  $50\%VO_2 = (.1\text{mL/kg/min} \cdot 80.4\text{m/min}) + (80.4\text{m/min} \cdot \%grade \cdot 1.8\text{mL/kg/min}) + (3.5\text{mL/kg/min})$ . The speed of 3 mph was

kept constant to ensure that all participants would maintain a walking gait pattern. It has been shown that running requires two times the amount of oxygen as walking (11).

**Table 1.** Run 12 kmh/7.5 mph VO<sub>2max</sub> test.

Time (min)	Speed (mph)	Grade (%)
1:00	4.4	0
2:00	5	0
3:00	5.6	0
4:00	6.2	0
5:00	6.9	0
6:00	7.5	0
7:00	7.5	2
8:00	7.5	4
9:00	7.5	6
10:00	7.5	8
11:00	7.5	10
12:00	7.5	12
13:00	7.5	14
14:00	7.5	16
15:00	7.5	18
16:00	7.5	20
17:00	7.5	22
18:00	7.5	24
19:00	7.5	25

Prior to exercising, the participants resting heart rate was palpated for fifteen seconds and multiplied by four to determine beats per minute. The subjects would lie prone for ten minutes prior to palpation to ensure a true rested state. The subjects then warmed up for three minutes at a speed of 3 mph and 0% grade. Following the three-minute warm-up the grade was increased to the calculated value (value expected to yield 50% of reserve) for 5 minutes. VO<sub>2</sub> and HR were monitored using the PARVO system. ACSM guidelines state that steady state is reached within 1-4 minutes during light exercise (10); consequently, a 5-minute exercise bout was used to ensure steady state was achieved. After the test, calculations were made to determine what

## VO<sub>2</sub> RESERVE VS. HR RESERVE

percent of reserves (VO<sub>2</sub>R and HRR) had been reached. These values were then compared to the pre-test 50% calculation.

### Statistical Analysis


While subjects performed the exercise bout at the predicted %50VO<sub>2</sub>R, VO<sub>2</sub> and HR were obtained on a minute-by-minute basis. The mean VO<sub>2</sub> (mL/kg/min) and HR (bpm) were then calculated for each subject between the fifth and eighth minute of exercise. Statistical analysis was carried out using IBM SPSS Statistics (Version 20). Paired *t*-tests were performed to determine the difference between the predicted and actual values for VO<sub>2</sub> and HR at 50% of their reserves. Correlational analysis via Pearson's *r* was used to compare the percent of VO<sub>2</sub>R and HRR achieved. Statistical significance was determined at  $p < 0.05$ , and Standard Error of Estimate (SEE) was also calculated.

## RESULTS


All results were reported as mean  $\pm$  standard deviation. Statistical analysis revealed that the predicted 50%VO<sub>2</sub> ( $27.1 \pm 3.3$ ) was significantly greater than the actual VO<sub>2</sub> achieved ( $25.6 \pm 3.3$ ),  $t(27) = 8.2$ ,  $p < .001$  (Figure 1). It was also found that the mean predicted HR at 50% of the reserve ( $128.8 \pm 7.5$ ) was significantly less than the actual HR achieved ( $135.6 \pm 10.6$ ),  $t(27) = 5.4$ ,  $p < .001$  (Figure 2).

The correlation between measured and predicted VO<sub>2</sub> at 50%R was  $r = .96$ ,  $p < .001$ , with an SEE = .94 mL/kg/min. The correlation between measured and predicted HR at 50%R was  $r = .78$ ,  $p < .001$ , with an SEE = 6.6 bpm.


In addition to *t*-tests, Pearson's *r* was used to analyze the correlation between %VO<sub>2</sub>R and %HRR during moderate intensity exercise. It was found that a statistically non-significant, weak positive, relationship existed between the two variables ( $r = .31$ ,  $p = .105$ ) (Figure 3).



**Figure 1.** Regression line of predicted and actual VO<sub>2</sub> @ 50%R. Shows that a strong correlation existed ( $r = .96$ ) and that predicted VO<sub>2</sub> was significantly greater ( $27.1 \pm 3.3$ ) than actual VO<sub>2</sub> achieved ( $25.6 \pm 3.3$ ),  $t(27) = 8.2$ ,  $p < .001$ .



**Figure 2.** Regression line of predicted and actual HR @ 50%R. Shows that a moderate correlation existed ( $r = .78$ ) and that predicted HR was significantly less ( $128.8 \pm 7.5$ ) than actual HR achieved ( $135.6 \pm 10.6$ ),  $t(27) = 5.430$ ,  $p < .001$ .



**Figure 3.** Regression line for %HRR at a given %VO<sub>2</sub>R. A statistically non-significant, weak positive, relationship existed between the two variables ( $r = .31$ ,  $p = .105$ ).

## DISCUSSION

The primary purpose of this study was to determine the relationship between VO<sub>2</sub>R and HRR during moderate intensity exercise. Secondly, determining the predictability of VO<sub>2</sub> and HR at moderate intensity exercise. It was hypothesized that VO<sub>2</sub>R and HRR would be directly correlated at moderate intensity exercise (50%R). It was also hypothesized that VO<sub>2</sub> and HR could accurately be predicted. This study indicated that VO<sub>2</sub> could be more accurately predicted than HR during moderate intensity exercise. Also, a weak positive correlation existed between VO<sub>2</sub>R and HRR. This indicated that it was less reliable to predict one variable given another (i.e., HR cannot be accurately predicted when the VO<sub>2</sub> of a given exercise bout is known). On the basis of a 50%R (moderate intensity) prediction, mean %VO<sub>2</sub>R values were over-predicted (Actual %VO<sub>2</sub>R = 46.9% ± 2.0%) whereas mean %HRR values were under-predicted (Actual %HRR = 55.3% ± 5.4%).

Our results on the prediction of VO<sub>2</sub> supported the findings of Cunha et al., which examined the association between VO<sub>2</sub>R and HRR as the intensity, and duration, of exercise were increased (28 physically active males). Cunha's study concluded that predicted VO<sub>2</sub> values were significantly greater than actual VO<sub>2</sub> values. Additionally, it was found that as exercise intensity increased the strength of the relationship between %VO<sub>2</sub>R and %HRR weakened (3).

The results of our study were shown to overpredict VO<sub>2</sub>, however, the SEE value of .94 mL/kg/min suggests that accurate predictions may still be made in most practical circumstances. This finding supports the research conducted on 26 highly trained, healthy male cyclists by Lounana et al. They found that VO<sub>2</sub> could be accurately predicted when using reserve values (5).

However, the results of this study that were dependent on HR data deviated from the majority of previous research findings. The majority of past research had found a direct correlation between %VO<sub>2</sub>R and %HRR (i.e., enabling accurate predictions of HR from VO<sub>2</sub>). Swain et al. investigated the relationship between %VO<sub>2</sub>R and %HRR in apparently healthy individuals (26 males, 24 women, 18-20 yrs) during treadmill exercise they found that the regression between VO<sub>2</sub>R and %HRR was  $r = 0.99 \pm 0.002$  (7) whereas our results found a weak correlation ( $r = .31$ ).

These differences could have been the result of many different factors, the first of which is elevated levels of catecholamines in the blood circulating throughout the

body. Catecholamines are hormones released by the body during exercise and are responsible for elevating blood pressure, breathing rate, and heart rate (6). Many participants in this study had exercised prior to performing the submaximal portion of the study. This would have resulted in elevated levels of these hormones. It was assumed that by lying prone for 10 minutes, an accurate resting HR, as well as an accurate ambulatory HR would be achieved. However, it is plausible that catecholamines were still circulating through the system, causing the participants resting HR to be elevated above that which was predicted.

Some other limitations consisted of making the assumption of 3.5mL/kg/min as the value for resting VO<sub>2</sub> as well as inconsistency of HR measurements. The value of 3.5mL/kg/min was assumed for all participants as resting VO<sub>2</sub> but upon further research it was found that resting VO<sub>2</sub> has been shown to vary greatly from subject to subject. Cunha et al. found that their subjects measured resting VO<sub>2</sub> values were significantly greater than the assumed value of 3.5mL/kg/min ( $4.3 \pm 0.9$  mL/kg/min,  $p < .001$ ) (3). Conversely, in a study conducted by Byrne et al. that looked at two separate cohorts (593 women, 78 men, 18-74 yrs; 49 women 49 men,  $38 \pm 5$  yr) resting VO<sub>2</sub> values were found to be significantly lower than 3.5mL/kg/min ( $2.6 \pm 0.4$  mL/kg/min) (2).

When collecting exercising HR data, a HR monitor was used. However, resting HR's were palpated, using the radial pulse, for fifteen seconds and multiplied by four. This method, while convenient and easy to use, has a large margin for error. The counting

of one extra or one less beat would result in a difference of  $\pm 4$  beats per minute. By using a HR monitor to obtain both resting and exercising HR's better consistency in readings would have been possible. A more meticulous approach to the collection of data could potentially indicate that HR can be accurately predicted using reserves. Furthermore, better collection of HR data could also indicate that a direct relationship between %VO<sub>2</sub>R and %HRR does exist. Such results would support the findings of Brawner and associates, Dalleck and Kravitz, Suain and Leutholtz, and Swain and colleagues (1,4,8,9). Lastly, it is possible that not all participants reached the desired exercise intensity, as the workload was a predictive calculation derived from their maximal test.

In conclusion, this study indicated that VO<sub>2</sub> could be more accurately predicted than HR during moderate intensity exercise. The weak correlation between VO<sub>2</sub>R and HRR indicates that caution should be used when relying on a HR to determine VO<sub>2</sub>.

## REFERENCES

1. Brawner CA, Keteyian SJ, Ehrman JK. The relationship of heart rate reserve to VO<sub>2</sub> reserve in patients with heart disease. *Med Sci Sports Exerc* 34(3): 418-422, 2002.
2. Byrne NM, Hills AP, Hunter GR, Weinsier RL, Schutz Y. Metabolic equivalent: one size does not fit all. *J Appl Physiol* 99: 1112-1119, 2005.
3. Cunha FA, Midgley AW, Monteiro WD, Campos FK, Farinatti PTV. The relationship between oxygen uptake reserve and heart rate reserve is affected by intensity and duration during aerobic exercise at constant work rate. *Appl Physiol Nutr Metab* 36(6): 839-847, 2011.

## VO2 RESERVE VS. HR RESERVE

4. Dalleck LC, Kravitz L. Relationship between % heart rate reserve and % VO<sub>2</sub> reserve during elliptical crosstrainer exercise. *J Sports Sci and Med* 5: 662-671, 2006.
5. Lounana J, Campion F, Noakes TD, Medelli J. Relationship between %HRmax, %HR Reserve, %VO<sub>2max</sub>, and %VO<sub>2</sub> Reserve in Elite Cyclists. *Med Sci Sport Exer* 39(2): 350-357, 2007.
6. Schwarz NA, Spillane M, Bounty PL, Grandjean PW, Leutholtz B, Willoughby DS. Capsaicin and evodiamine ingestion does not augment energy expenditure and fat oxidation at rest or after moderately-intense exercise. *Nutr Res* 33: 1034-1042, 2013.
7. Swain DP, Leutholtz BC. Heart rate reserve is equivalent to % VO<sub>2</sub>Reserve, not to %VO<sub>2max</sub>. *Med Sci Sports and Exer* 29(3): 410-414, 1997.
8. Swain DP, Leutholtz BC, King ME, Haas LA, Branch DJ. Relationship between % heart rate reserve and % VO<sub>2</sub> reserve in treadmill exercise. *Med Sci Sports Exer* 30(2): 318-321, 1998.
9. Thompson WR, Gordon NF, Pescatello LS. *ACSM's Guidelines for Exercise Testing and Prescription*. 8<sup>th</sup> ed. Philadelphia (Pa): Lippincott Williams & Wilkins; 2010. 453 p.
10. Thompson WR, Gordon NF, Pescatello LS. *ACSM's Guidelines for Exercise Testing and Prescription*. 8<sup>th</sup> ed. Philadelphia (Pa): Lippincott Williams & Wilkins; 2010. 49 p.
11. Thompson WR, Gordon NF, Pescatello LS. *ACSM's Guidelines for Exercise Testing and Prescription*. 8<sup>th</sup> ed. Philadelphia (Pa): Lippincott Williams & Wilkins; 2010. 458 p.