

## **Low-density lipoproteins, body mass index, and the female sex are predictors of reduced cutaneous reactive hyperemia in human skin**

Craighead, D. Dahmus, J. Kenney, WL, FACSM. Alexander-Holowatz, L, FACSM. Penn State University, University Park, PA

[dhc139@psu.edu](mailto:dhc139@psu.edu), [jdd5081@psu.edu](mailto:jdd5081@psu.edu), [lma191@psu.edu](mailto:lma191@psu.edu), [w7k@psu.edu](mailto:w7k@psu.edu)

**Purpose:** The human cutaneous circulation is a representative vascular bed to assess systemic microvascular dysfunction associated with preclinical cardiovascular diseases. Laser speckle contrast imaging (LCSI) is a state-of-the-art method used to measure skin blood flow (SkBF), imaging a larger surface area compared to single-point laser-Doppler flowmetry. Using LCSI, we sought to determine what cardiovascular risk factors were most predictive of the cutaneous reactive hyperemic (RH) response in human subject with a broad range of age cardiovascular disease risk factors. **Methods:** A standardized RH protocol consisting of three, five minute suprasystolic occlusions was performed on 38 human subjects ranging in age (19-82 years), body mass index (BMI: 19.7-36.2 kg/m<sup>2</sup>), mean arterial pressure (MAP: 67-99 mmHg) and serum low-density lipoproteins (LDL) (54-185 mg/dl). Following the occlusion, the RH SkBF response was imaged using the full-field LCSI. Regions of interests (ROI) were determined, integrated, and normalized to cutaneous vascular conductance (CVC, integrated ROI flux/MAP). A stepwise regression analysis was performed to determine the best predictors of RH parameters including the peak SkBF, and the total hyperemic response (THR). **Results:** Subject sex ( $R^2=0.158$ ,  $P<0.001$ ), BMI ( $R^2=0.195$ ,  $P<0.001$ ), and serum LDL ( $R^2=0.27$ ,  $P<0.001$ ) were all significant predictors of peak RH CVC. Both age ( $R^2=0.108$ ,  $P<0.001$ ) and BMI ( $R^2=0.147$ ,  $P=0.012$ ) significantly predicted the THR. **Conclusion:** Increased BMI, serum LDL, and the female sex, are associated with decreased peak SkBF response during RH. Increased age and BMI are associated with a decreased total hyperemic response. (NIH R01 AG07004-19-25)