

## High Dietary Sodium Reduces Flow Mediated Dilation in Humans with Salt Sensitive & Salt Resistant BP.

Matthews, E.L., DuPont, J.J., Greaney, J.L., Lennon-Edwards, S.L., Edwards, D.G., Farquhar, W.B. University of Delaware, Newark, DE

[elmatthe@udel.edu](mailto:elmatthe@udel.edu), [jdupont@udel.edu](mailto:jdupont@udel.edu), [jgreaney@udel.edu](mailto:jgreaney@udel.edu), [slennon@udel.edu](mailto:slennon@udel.edu), [dge@udel.edu](mailto:dge@udel.edu), [wbf@udel.edu](mailto:wbf@udel.edu)

Human and animal studies have demonstrated that dietary sodium loading impairs endothelial function. Recent data suggests that this impairment may occur independent of a change in blood pressure (BP) (i.e., in those with 'salt resistant' BP). However, a sodium-induced increase in BP (i.e., 'salt sensitive' BP) may amplify the negative effects of sodium on the endothelium. **Purpose:** We tested the hypothesis that dietary sodium loading would impair flow mediated dilation (FMD) more in those with salt sensitive (SS) BP compared to those with salt-resistant (SR) BP. **Methods:** Five SS (2 men, 3 women; age  $43 \pm 4$  years) and 7 SR (3 men, 4 women; age  $45 \pm 3$  years) subjects were enrolled in a controlled feeding study where all food was provided. The diet consisted of a run-in period (3-7 days, 100mmol sodium/day) immediately followed by a two phase randomized crossover seven-day diet perturbation: low sodium (LS); 20 mmol/day and high sodium (HS); 300-350 mmol/day. FMD of the brachial artery was assessed on the last day of each diet condition. Mean arterial BP (MAP) was assessed using a 24-hr ambulatory monitor. SS BP was defined as a change in MAP of  $> 5$  mmHg between the low and the high sodium diets. **Results:** By study design, MAP increased in the SS group during the HS condition (LS:  $83.0 \pm 1.5$ , HS  $90.2 \pm 1.7$  mmHg,  $p < 0.05$ ), but not the SR group (LS:  $83.6 \pm 3.0$ , HS  $83.1 \pm 3.1$  mmHg,  $p > 0.05$ ). HS attenuated FMD in both the SS group (LS:  $10.5 \pm 2.1\%$ , HS  $6.9 \pm 1.3\%$ ,  $p < 0.05$ ) and SR group (LS:  $12.8 \pm 2.0\%$ , HS  $8.2 \pm 1.2\%$ ,  $p < 0.05$ ) with no difference between groups for the change in FMD (SS:  $\Delta -3.6 \pm 1.4\%$ , SR:  $\Delta -4.6 \pm 1.0\%$ ,  $p > 0.05$ ). **Conclusion:** These preliminary data indicate that a high sodium diet impairs FMD to a similar extent in SS and SR individuals, suggesting that the effects of dietary sodium on vascular endothelial function may be independent of BP.

Research funded by NIH grants 2 P20 RR016472-11 and R01 HL104106.

---