

The Effects of Physical Activity on Markers of Hepatic Inflammation During Weight-Loss

Seija A, Poole KE, Eck K, Harris M, Nick TN, and Wooten JS

Exercise Biochemistry and Physiology Laboratory; Department of Kinesiology and Health Education; Southern Illinois University Edwardsville; Edwardsville, IL

Category: Undergraduate

Advisor / Mentor: Wooten JS (jwooten@siue.edu)

ABSTRACT

Non-alcoholic fatty liver disease (NAFLD) represents a continuum that begins with accumulation of lipid in hepatic cells progressing to hepatic steatosis with inflammation (steatohepatitis), fibrosis, and cirrhosis. Weight-loss using dietary modification and physical activity are common strategies used for the treatment of NAFLD; however, it remains to be determined the effects of physical activity on hepatic inflammation during weight-loss. The purpose of this study was to determine the therapeutic role of physical activity on plasma and hepatic inflammatory markers during weight-loss. Male C57BL/6 mice were fed either a low-fat (LFD; 10% kcal fat) or high-fat (HFD; 60% kcal fat) diet for 10-weeks. Following 10-weeks, the HFD group was randomly assigned to either a LFD (Diet) or LFD with physical activity (Diet+PA) to induce weight loss for 8-weeks. After 8-weeks, reductions in body mass were observed in both Diet and Diet+PA groups (see Table 1.). Interestingly, the Diet+PA group lost significantly ($P<0.05$) more body mass than the Diet group. Despite significant ($P<0.05$) reductions in body mass and HOMA-IR, plasma TNF- α remained elevated in the Diet and Diet+PA groups. Moreover, Diet+PA plasma TNF- α was significantly ($P<0.05$) greater than the HFD obese controls. Elevated plasma TNF- α in the Diet+PA was matched by a greater hepatic expression of IL-1 β and IL-6 mRNA when compared to all groups. Interestingly, the expression of TGF- β 1 mRNA was significantly ($P<0.05$) reduced in the Diet+PA when compared to all groups. The elevated plasma TNF- α and expression of IL-1 β and IL-6 mRNA are likely due to physical activity. It remains unclear as to the pro-inflammatory effects of physical activity during weight-loss; however, this may be part of a protective adaption to regular exercise. Furthermore, the reduced hepatic TGF- β 1 mRNA levels suggest a protective strategy against fibrogenesis in the spectrum of liver disease.

Table 1. Whole body and hepatic metabolic characteristics following weight-loss.

Variables	LFD (n=12)	HFD (n=12)	Diet (n=12)	Diet+PA (n=12)
Body mass (g)	30.2 \pm 1.1	48.8 \pm 0.5*	30.3 \pm 0.7 [†]	26.1 \pm 0.3* ^{††}
HOMA-IR	22.9 \pm 1.2	187.3 \pm 7.5*	19.4 \pm 8.8 [†]	25.3 \pm 10.5 [†]
IL-6 (pg/mL)	6.4 \pm 0.7	6.2 \pm 1.0	5.9 \pm 0.9	6.4 \pm 0.9
TNF- α (pg/mL)	30.8 \pm 6.7	60.6 \pm 5.3*	74.0 \pm 8.1*	82.5 \pm 7.7* [†]
IL-1 β mRNA	1.00 \pm 0.51	0.97 \pm 0.34	1.20 \pm 0.59	2.83 \pm 0.62* ^{††}
IL-6 mRNA	1.00 \pm 0.45	1.53 \pm 0.50	1.16 \pm 0.72	2.36 \pm 0.55* ^{††}
TNF- α mRNA	1.00 \pm 0.09	0.89 \pm 0.08	0.94 \pm 0.14	0.83 \pm 0.06
TGF- β 1 mRNA	1.00 \pm 0.06	1.02 \pm 0.06	1.02 \pm 0.10	0.84 \pm 0.05 [†]

Note. Data are presented as mean \pm SEM. *Significantly ($P<0.05$) different than LFD; [†]significantly ($P<0.05$) different than HFD; ^{††}significantly ($P<0.05$) different than Diet.