



Original Research

Short Term High-Repetition Back Squat Protocol Does Not Improve 5-km Run Performance

MATTHEW J. BARENIE^{†1,5}, JORDAN T. DOMENICK^{†1}, JASON E. BENNETT^{‡2}, GEORGE G. SCHWEITZER^{‡3}, PAULINA SHETTY^{‡4}, and EDWARD P. WEISS^{†1}

¹Department of Nutrition and Dietetics, Saint Louis University, St. Louis, MO, USA; ²Department of Physical Therapy, Carroll University, Waukesha, WI, USA; ³Department of Medicine, Washington University School of Medicine, St. Louis, MO, USA; ⁴Science Department, Johnson County Community College, Overland Park, KS, USA; ⁵Human Performance and Exercise Biochemistry Laboratory, Department of Kinesiology, Indiana University, Bloomington, IN, USA

[†]Denotes graduate student author, [‡]Denotes professional author

ABSTRACT

International Journal of Exercise Science 13(7): 1770-1782, 2020. The purpose of this study was to evaluate the hypothesis that a novel high-repetition, low-resistance back squat training protocol, designed to stimulate high-intensity interval training, improves 5-km run performance. Fifteen runners [4 male, 11 female; 150 + minutes of endurance exercise/week; age = 22.7 ± 2.0 y; 21.5 ± 2.2 kg/m² BMI] in this single-group test-retest design completed two weeks of back squats consisting of three sets of 15-24 repetitions at 60% of estimated one-repetition max (1RM), three times per week (1-2 days of rest between sessions). Outcome tests included a 5-km outdoor timed run, laboratory indirect calorimetry to quantify substrate oxidation rates during steady-state submaximal exercise (60% and 70% heart rate max (HRmax)), and estimated 1RM for back squats. Back squat estimated 1RM increased by 20% (58.3 ± 18.5 to 70.2 ± 16.7 kg, *P* < 0.001). However, 5-km run times due to the back squat protocol did not significantly change (Pre-Squats: 23.9 ± 5.0 vs. Post-Squats: 23.7 ± 4.3 minutes, *P* = 0.71). Likewise, the squat training program did not significantly alter carbohydrate or lipid oxidation rates during steady-state submaximal exercise at 60% or 70% of HRmax (*P* values ranged from 0.36 - 0.99). Short term high-repetition back squat training does not appear to impact 5-km run performance or substrate utilization during submaximal exercise.

KEY WORDS: Barbell training, lower body resistance exercise, long-distance running, carbohydrate and fat utilization

INTRODUCTION

Intense, sprint-type anaerobic exercise (commonly referred to as high-intensity interval training, HIIT) improves aerobic exercise capacity (9,17-19). The nature of the “sprints” used in these protocols vary but in general consist of 30 second intervals of maximal effort cycling,

interspersed by recovery periods lasting three minutes (9,17,18). The improvements in endurance performance seen in recreationally active individuals after HIIT training appear to be at least partly attributable to increases in muscle content of oxidative mitochondrial enzymes (8,18,30), including signaling proteins for mitochondrial biogenesis (3,29), enhanced muscle buffering capacity (17,33), and increased glycogen stores (17). These improvements in performance and adaptations in muscle also appear to occur in as little as six sessions over two weeks (9,17) and signals for mitochondrial biogenesis increasing after just one HIIT session (29,43).

In many respects, back squat strength training may impose similar physiologic demands on the body as anaerobic sprint cycle training, especially from a metabolic perspective (6). Further, back squats have been shown to be safe as well as effective in preventing against lower body injury and significantly increasing lower body strength (14,24,36). Programs designed to improve endurance performance often center on increases in training load, and subsequently expose athletes to an increased risk of injury (12,45). Running injury prevalence has been shown to affect ~20-80% of runners (46). Methods to improve exercise performance while minimizing injury risk (e.g., back squatting) would be highly desirable.

A typical set of strength training, consisting of 8 – 12 repetitions with a load equivalent to 70-80% of 1RM causes muscle failure in approximately 30 seconds (28). With somewhat lighter loads and more repetitions, a set of strength training exercise can last 60 seconds or even longer. In this timeframe both HIIT training and high-repetition strength training put high demands on ATP-PC (alactic) and anaerobic glycolysis (lactic) energy systems. Based on these likely similarities, it is plausible that high-repetition back squat training could have similar effects as HIIT training, in that it might improve endurance performance. Further, HIIT has been shown to increase the capacity for skeletal muscle lipid oxidation and work output during sub-maximal exercise, suggestive of a glycogen sparing effect (2,8). Considering the benefits and similarities described above regarding strength training and HIIT, no studies to date have examined the effects of a short term back squat protocol on 5-km running performance. As evidenced by Hurst and Board, the 5-km outdoor run time trial is a reproducible exercise performance test which is indicated by an intraclass correlation coefficient of 0.984 (25).

The primary purpose of this study is to evaluate the effects of two weeks of a novel high-repetition back squat exercise on 5-km running performance in recreational endurance athletes. A secondary purpose of this study was to assess the effects of the squat protocol on substrate utilization in endurance athletes. If the protocol is found beneficial, high-repetition back squat exercise training could be an effective strategy for runners, and possibly other endurance athletes, to improve their performance.

METHODS

Participants

A power analysis conducted with G*POWER 3.1.5 (University of Kiel, University of Dusseldorf, and University of Mannheim, Germany) determined that 15 participants were needed in the

present study for a power of 0.80, with an effect size of 0.78 and an $\alpha = 0.05$. Healthy individuals aged 18-30 years were recruited from the St. Louis, Missouri metropolitan area. Volunteers completed a medical history and medications questionnaire, which was used along with criteria from the American College of Sports Medicine to classify each individual as low, moderate, or high risk for medical complications during exercise (34). Both moderate- and high- risk individuals were excluded. Subjects were required to be recreational or higher-level endurance athletes who performed at least 150 minutes of endurance exercise per week. There was no specific 5-km completion time necessary to participate in this study. Subjects also must not have performed lower body resistance training within six weeks of beginning the study. The average subject characteristics correspond to a normal weight body mass index ($21.5 \pm 2.2 \text{ kg/m}^2$), young adult (age = 22.7 ± 2.0 years), with a female to male ratio of 11:4. For the female, average height was $164.5 \pm 6.7 \text{ cm}$ and weight was $56.1 \pm 8.5 \text{ kg}$. For men, average height was $177.8 \pm 9.9 \text{ cm}$ and weight was $72.8 \pm 12.8 \text{ kg}$. All participants provided written informed consent prior to participation and the Saint Louis University Institutional Review Board for the protection of human subjects approved the protocol. This research was carried out fully in accordance to the ethical standards of the International Journal of Exercise Science (35).

Protocol

Study design: This study was an uncontrolled single-group test-retest design, in which subjects underwent outcomes assessments before and after the 2-week squat exercise training protocol. The outcome tests were conducted in the following order: 5-km run, assessment of substrate oxidation rates during exercise, and finally estimated one-repetition max (1RM). There was approximately 15-20 minutes between the 5-km and substrate oxidation assessments and approximately five minutes between substrate oxidation measures and estimated 1RM.

Training intervention: The squat training protocol consisted of six sessions of high-repetition barbell back squats over two weeks with 1 - 2 days of rest in between sessions. The participants were required to perform three sets of squats (end range-of-motion was defined as thighs parallel to the floor) per session with approximately two minutes rest between sets. The initial load for the squats was set at 60% of 1RM, which was determined at the initial assessment by means of the Berger Prediction Table (4). The load was either increased or decreased by 2.3 - 4.5 kilograms as needed to allow subjects to complete 15-24 repetitions in each set of squats. The subjects were encouraged to use maximal or near-maximal effort during every set (e.g., continue exercise until volitional fatigue or within one or two repetitions of volitional fatigue, as has been used in other studies) (18,41). To increase safety and consistency, a power rack equipped with safety bars was utilized and each squat training session was supervised by American College of Sports Medicine Certified Personal Trainers.

Endurance exercise performance: The endurance exercise performance test was a 5-km run that took place using the Saint Louis University's outdoor 400-meter track (12.5 laps). Run time information was not provided and participants were not allowed to time themselves but were informed to complete each time trial in the least amount of time possible. Participants were equipped with Polar FT1 heart rate monitors (Bethpage, NY) and immediately after run completion, heart rate and rating of perceived exertion were obtained (7).

Exercise metabolism: Indirect calorimetry (TrueOne 2400; ParvoMedics, Murray, UT) was used to measure oxygen consumption (VO₂) and carbon dioxide production (VCO₂) and respiratory exchange ratio (RER) was calculated (VCO₂/VO₂). Carbohydrate and fat oxidation rates were calculated based on VO₂ and VCO₂ according to the following equations (10):

$$\text{Cho (g/min)} = -2.909 * \text{VO}_2 \text{ (L/min)} + 4.115 * \text{VCO}_2 \text{ (L/min)} - 2.539 * \text{Uun}$$

$$\text{Fat (g/min)} = 1.689 * \text{VO}_2 \text{ (L/min)} - 1.689 * \text{VCO}_2 \text{ (L/min)} - 1.943 * \text{Uun}$$

Uun (urine nitrogen as estimate of protein utilization) was assumed to be zero

Breath-by-breath metabolic measures were performed during steady-state submaximal exercise on a treadmill. Prior to each test, a 3-liter air syringe (Hans Rudolph, Inc., Shawnee, KS) was used to generate a series of flow profiles for calibrating the pneumotach flow meter on the calorimeter. The carbon dioxide and oxygen analyzers were calibrated prior to each test with medical grade gases of known carbon dioxide and oxygen concentrations. The initial treadmill speed was self-selected by the participant usually as a slow walking speed, while the grade on the treadmill was 0% throughout the measure. The speed was subsequently increased by 0.3 - 1.0 mph every two minutes until ~60% heart rate max (HR_{max}) was achieved. Heart rate goals were determined by calculating age predicted HR_{max}, using the formula $208 - (0.7 \times \text{age})(42)$. Once heart rate in the ~60% range was achieved, the participant was held at the corresponding speed for 10 minutes. Following, speed was increased in stages until ~70% HR_{max} was attained and again, the speed was then held constant for 10 minutes. The treadmill speeds in the pre-squat training assessment were then replicated in the post-squat training assessment. Metabolic data averaged over 60 second intervals from the last five minutes of each stage was then averaged to reflect steady-state metabolic responses to exercise.

Exercise and dietary control: Participants were advised to maintain their usual regimen of endurance training throughout their participation in the study and were instructed to continue to not engage in any lower body strength exercise. The participants kept a food and beverage journal and an exercise activity log for two days prior to the pre-squat training testing session and then used the recorded information to follow a similar diet and activity pattern leading up to the post-squat training testing sessions. Researchers photocopied the subject's diet and exercise log and distributed them to each subject two days before post-squat training testing. Each participant was continually encouraged to follow their respective regular diet and exercise routine during the entire duration of this study.

Ambient weather conditions: Ambient air temperature, relative humidity, and heat index at the time of each run were acquired from the National Weather Service (www.weather.gov).

Statistical Analysis

Data were summarized as means \pm standard deviation. The mean difference scores (e.g. final minus baseline values) were tested to determine if they were significantly different from zero

using a paired t-test. Regression analysis were performed to identify predictive relationships among variables. A P -value of < 0.05 was considered significant. Analyses were performed with Microsoft Excel and IBM SPSS Statistics (version 24) software.

RESULTS

Exercise training compliance and muscle strength: All subjects completed the required assessments and squat exercise training sessions according to protocol (100% compliance). Participants verbally expressed their understanding and willingness to adhere to their usual exercise and dietary regimens throughout the study. As evidence of compliance with the training program, and also of the efficiency of the short term back squat training protocol on muscle strength, estimated 1RM for the squat exercise increased by 20%, from 58.3 ± 18.5 to 70.2 ± 16.7 kg ($P < 0.001$) during the training period.

Exercise performance: Five-kilometer run performance did not change in response to squat training ($P = 0.71$). During the post-squat training run, heart rate, as measured immediately after the 5-km run, was 6% higher ($P = 0.062$; Table 1), albeit not quite significantly, and RPE was 8% higher ($P = 0.048$). As depicted in Table 2, ambient air temperature ($P = 0.007$) and heat index ($P = 0.006$) were significantly higher during the post-squat training run; relative humidity did not differ between trials ($P = 0.42$). Because high heat stress may negatively affect exercise performance (20,26,31,38,44,47), we performed a regression analysis to determine if changes in heat stress were predictive of the changes in 5-km run times. However, neither the change in air temperature between trials ($P = 0.29$) or the difference in heat stress ($P = 0.54$) were predictive of the changes in running performance, suggesting that changes in weather did not affect the results. We also used regression analysis to determine if baseline fitness (as approximated by using the pre-squat training run time) was predictive of the changes in running performance. Indeed, it was predictive (beta-coefficient: -0.20 ± 0.09 , $P = 0.04$, r-square: 0.28), indicating that runners with slower baseline run times had greater improvements in performance in response to squat training. In light of this finding, we performed a sub-analysis after excluding the slowest runners (> 24 minute baseline 5-km time, $n = 6$); the analysis of 5-km results remained non-significant ($P = 0.17$). Likewise, we analyzed the results after excluding the fasted participants (< 21 minute baseline 5-km run time, $n = 5$) and the results remained non-significant ($P = 0.68$). Thus, while the magnitude of change in run times appears to partly depend on baseline fitness, it does not appear that either slow or fast runners could have cancelled out effects in the remainder of the sample. Figure 1 shows the individual responses of the runners for 5-km run times.

Table 1. Effect of high repetition back squat training on 5-km run performance, heart rate, and rating of perceived exertion.

	Pre-Squats	Post-Squats	Delta Scores	P-value
5-km Run Time, minutes	23.9 ± 5.0	23.7 ± 4.3	-0.2 ± 1.9	0.71
Heart Rate immediate post exercise, BPM	174.9 ± 18.8	186.2 ± 10.4	11.3 ± 21.5	0.062
RPE at end of 5-km run, 6-20 scale	14.0 ± 2.1	15.1 ± 1.5	1.1 ± 1.9	0.048

Values are means ± standard deviations. Delta scores were calculated as follow-up value minus baseline value. P-values reflect the significance of T-tests which compared baseline and follow-up values. BPM, beats per minute. RPE, rating of perceived exertion.

Table 2. Ambient weather conditions during the outdoor 5-km performance runs.

	Pre-Squats	Post-Squats	P-value
Temperature, °C	24.6 ± 3.6	28.7 ± 4.2	0.007
Relative Humidity, %	62.1 ± 11.6	66.9 ± 18.8	0.42
Heat Index, °C	25.0 ± 4.2	31.8 ± 7.5	0.006

Values are means ± standard deviations. P-values reflect the significance of T-tests which compared baseline and follow-up values. National Weather Service data were acquired from www.weather.gov.




Figure 1. Participant-level 5-km run times from before and after the two week high-repetition back squat training period. Lines represent the changes from pre- to post-squat training for each participant. The solid bars represent the means and the associated values reflect means ± standard deviations. The 5-km runs were performed on an outdoor 400-meter running track. The P-value reflects the statistical significance of the difference between trials.

Table 3. Effect of high repetition back squat training on metabolic outcomes during steady-state submaximal exercise.

	Pre-Squats	Post-Squats	P-value
Respiratory Exchange Ratio			
60%	0.80 ± 0.05	0.80 ± 0.06	0.98
70%	0.85 ± 0.05	0.85 ± 0.04	0.93
Oxygen Consumption, ml/kg/min			
60%	12.7 ± 5.4	12.8 ± 5.2	0.94
70%	20.3 ± 4.4	21.0 ± 4.7	0.36
Carbohydrate Oxidation, % of EE			
60%	35 ± 17	37 ± 19	0.74
70%	49 ± 16	49 ± 15	0.97
Carbohydrate Oxidation, kcal/min			
60%	1.9 ± 1.6	1.4 ± 1.1	0.42
70%	3.1 ± 1.6	3.2 ± 1.5	0.77
Fat Oxidation, % of EE			
60%	65 ± 16	63 ± 18	0.74
70%	51 ± 16	51 ± 15	0.96
Fat Oxidation, kcal/min			
60%	2.4 ± 1.0	2.4 ± 1.1	0.99
70%	3.0 ± 0.9	3.1 ± 0.9	0.66
Total Energy Expenditure, kcal/min			
60%	3.7 ± 1.8	3.8 ± 1.7	0.86
70%	6.0 ± 1.8	6.2 ± 1.9	0.38
Heart Rate, BPM			
60%	116.1 ± 2.5	115.7 ± 1.7	0.55
70%	135.4 ± 2.9	135.5 ± 1.7	0.83
Heart Rate, % of age-predicted max			
60%	61 ± 2	60 ± 1	0.54
70%	71 ± 1	71 ± 1	0.83

Values are means ± standard deviations. *P*-values reflect the significance of T-tests which compared baseline and follow-up values. EE = energy expenditure. 60% and 70% indicate percentages of age-predicted maximum heart rate.

Submaximal exercise substrate oxidation: During the lower steady-state work rate, percentage of HRmax was equivalent to 61 ± 2% at pre-squats and 60 ± 1% during post-squats testing. At the higher steady-state work rate, percentage of HRmax was equivalent to 71 ± 1% for both conditions. Respiratory exchange ratio at 60% of HRmax at pre-squats was 0.80 ± 0.05 and 0.80 ± 0.06 (*P* = 0.98) during post-squat testing. RER at 70% of HRmax was found pre-squats of 0.85 ± 0.05 and post-squats of 0.85 ± 0.04 (*P* = 0.93). In accordance with these findings, and because substrate oxidation rates are calculated from RER values, neither lipid oxidation rates nor

carbohydrate oxidation rates (at 60% and 70% of HRmax) were affected by the squat-training program. Further, oxygen consumption (ml/kg/min) at both steady-state work rates was not significantly altered by the training program. Energy expenditure (kcal/min) was found to not be affected by the squat protocol in either steady-state work rate. Table 3 shows all the results from the indirect calorimetry testing, none of which revealed significant differences from pre-squat to post-squat training.

DISCUSSION

The purpose of this study was to determine if a novel, short term, high-repetition back squat training intervention yields adaptations that would be characteristic of endurance training. In particular, we proposed that high-repetition back squat training would increase endurance exercise performance, as reflected by 5-km run performance and that it would shift substrate selection during exercise to increase lipid oxidation rate and decrease carbohydrate oxidation. Contrary to our expectations, two weeks of high-repetition back squat training did not alter 5-km run time (pre-squats: 23 minutes and 54 seconds vs. post-squats: 23 minutes and 42 seconds) or substrate selection during steady-state exercise.

Various meta-analyses in the literature overwhelming support the use of HIIT to improve endurance performance (19,40), and in light of the similarities between high-repetition back squat exercise and sprint-interval training (in terms of the duration and intensity of effort during intervals), it's not clear why high-repetition back squat training failed to improve endurance exercise performance. However, several possibilities may be considered. It is possible that the squat training did not improve performance because the study participants may have already been performing some type of interval training as part of their endurance training routine (this was not assessed). If this were the case, they would have already been receiving a training stimulus from their interval training and the squat training might not have had additive effects. Another possible explanation, albeit purely speculative, is that the high-velocity muscle contractions that occur during sprint-interval training are critical for triggering improvements in aerobic metabolic capacity and improving endurance performance (muscle contraction velocity during squat exercise is slow by comparison). However, to our knowledge, no studies have compared metabolic adaptations in muscle that has been trained with the same work rate and differing contraction velocities.

Other explanations for the null findings may relate to the relationship between myocellular adaptations and endurance performance. Although the present study was not designed to evaluate mechanisms, some commentary about mechanisms is warranted. In regards to citrate synthase, not all data shows an increase in its activity as a result of HIIT (39,48). Further, some studies show no significant associations between oxidative enzyme activity and exercise performance (11,15). Beneficial effects such as enhanced PGC-1 α and mitochondrial biogenesis are a result of HIIT, but the main determinants of run performance is maximal oxygen uptake (VO_{2max}), fractional utilization of VO_{2max} and running economy (5). A recent systematic review highlights that strength training has no impact on VO_{2max} (3). In addition, this study's strength training protocol may have increased the subjects' body mass, reduced mitochondrial volume

density and capillary density, ultimately lowering relative $\text{VO}_{2\text{max}}$ (27). Further, the training intervention may not have been long enough to induce sufficient adaptations (1,22). However, several studies have demonstrated that high-intensity interval training for two weeks is sufficient for improving endurance performance (9,17,18). Lastly, since no differences were found in regards to oxygen consumption, macronutrient oxidation rates or RER, it is speculated that running economy was not beneficially altered due to the high-repetition back squat protocol and therefore 5-km run time was not affected.

While outdoor endurance performance testing has advantages in that it has greater relevance to real-world endurance performance than laboratory tests, the unpredictable nature of weather can result in potentially confounding effects on exercise performance. Indeed, a limitation in the present study was that heat stress was significantly greater during the post-squat training testing; as heat stress has been shown to reduce some types of endurance exercise performance (13,20,31,38), it is conceivable that the high heat stress may have masked improvements in endurance performance. However, our regression analysis did not identify the between-trial differences in ambient air temperature or heat index as important determinants of the changes in run times. This finding is in agreement with evidence from running performance data from world championship running events between 1999 and 2011 (23), in which 5-km running performance was found to be minimally affected by major differences in ambient temperature (in contrast, performance during longer distance running is greatly impaired, while sprint-type events are improved in high heat conditions). Taken together, our regression analysis paired with evidence from Guy et al. (23) suggest that the changes in heat stress during our study may not have confounded the results. Nonetheless, additional studies with better control of heat stress conditions are warranted. Other possible effects of the changes in heat stress in our study are evident. The trend toward higher heart rates and significant elevation of ratings of perceived exertion in the post-squat training trial are consistent with the notion that higher heart rates are needed to increase cutaneous blood flow for heat dissipation (21) and that heat stress increases effort perception (16,32,37), even when exercise intensity is unchanged.

In conclusion, two weeks of high-repetition back squats did not influence 5-km run performance or cause a shift in the body's use of fuel during submaximal exercise. At this point, short term high-repetition back squat training does not appear to provide benefits for endurance performance but is still advisable to recommend strength training for other purposes. The study did continue to prove the undeniable effects of back squats for increasing lower body strength, even in such a short time frame. These findings strongly suggest for further research on high-repetition back squats and endurance performance using a longer duration protocol.

ACKNOWLEDGEMENTS

We are extremely grateful to the study participants who took the time from their busy schedules to participate in the study. Without their participation, this study would not have been possible.

REFERENCES

1. Aagaard P, Andersen JL, Bennekou M, Larsson B, Olesen JL, Crameri R. Effects of resistance training on endurance capacity and muscle fiber composition in young top-level cyclists. *Scand J Med Sci Sports* 21(6): e298-307, 2011.
2. Barker AR, Day J, Smith A, Bond B, Williams CA. The influence of 2 weeks of low-volume high-intensity interval training on health outcomes in adolescent boys. *J Sports Sci* 32(8): 757-765, 2014.
3. Bartlett JD, Hwa Joo C, Jeong T-S, Louhelainen J, Cochran AJ, Gibala MJ. Matched work high-intensity interval and continuous running induce similar increases in PGC-1 α mRNA, AMPK, p38, and p53 phosphorylation in human skeletal muscle. *J Appl Physiol* 112(7): 1135-1143, 2012.
4. Berger RA. Determination of the resistance load for 1-RM and 10-RM. *J Assoc Phys Ment Rehabil* 15: 108-110, 1961.
5. Blagrove RC, Howatson G, Hayes PR. Effects of strength training on the physiological determinants of middle- and long-distance running performance: A systematic review. *Sports Med* 48(5): 1117-1149, 2018.
6. Bloomer RJ, Falvo MJ, Fry AC, Schilling BK, Smith WA, Moore CA. Oxidative stress response in trained men following repeated squats or sprints. *Med Sci Sports Exerc* 38(8): 1436-1442, 2006.
7. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc* 14(5): 377-381, 1982.
8. Burgomaster KA, Howarth KR, Phillips SM, Rakobowchuk M, MacDonald MJ, McGee SL. Similar metabolic adaptations during exercise after low volume sprint interval and traditional endurance training in humans. *J Physiol* 586(1): 151-160, 2008.
9. Burgomaster KA, Hughes SC, Heigenhauser GJ, Bradwell SN, Gibala MJ. Six sessions of sprint interval training increases muscle oxidative potential and cycle endurance capacity in humans. *J Appl Physiol* 98(6): 1985-1990, 2005.
10. Bursztein S, Glaser P, Trichet B, Taitelman U, Nedey R. Utilization of protein, carbohydrate, and fat in fasting and postabsorptive subjects. *Am J Clin Nutr* 33(5): 998-1001, 1980.
11. Costill DL, Fink WJ, Pollock ML. Muscle fiber composition and enzyme activities of elite distance runners. *Med Sci Sports* 8(2): 96-100, 1976.
12. Damsted C, Glad S, Nielsen RO, Sørensen H, Malisoux L. Is there evidence for an association between changes in training load and running-related injuries? *Int J Sports Phys Ther* 13(6): 931-942, 2018.
13. El Helou N, Tafflet M, Berthelot G, Tolaini J, Marc A, Guillaume M. Impact of environmental parameters on marathon running performance. *PloS one* 7(5): e37407, 2012.
14. Escamilla RF. Knee biomechanics of the dynamic squat exercise. *Med Sci Sports Exerc* 33(1): 127-141, 2001.
15. Foster C, Costill DL, Daniels JT, Fink WJ. Skeletal muscle enzyme activity, fiber composition and VO₂ max in relation to distance running performance. *Eur J Appl Physiol Occup Physiol* 39(2): 73-80, 1978.

16. Galloway SD, Maughan RJ. Effects of ambient temperature on the capacity to perform prolonged cycle exercise in man. *Med Sci Sports Exerc* 29(9): 1240–1249, 1997.
17. Gibala MJ, Little JP, Van Essen M, Wilkin GP, Burgomaster KA, Safdar A. Short-term sprint interval versus traditional endurance training: similar initial adaptations in human skeletal muscle and exercise performance. *J Physiol* 575(3): 901–911, 2006.
18. Gillen JB, Percival ME, Skelly LE, Martin BJ, Tan RB, Tarnopolsky MA. Three minutes of all-out intermittent exercise per week increases skeletal muscle oxidative capacity and improves cardiometabolic health. *PloS one* 9(11): e111489, 2014.
19. Gist NH, Fedewa MV, Dishman RK, Cureton KJ. sprint interval training effects on aerobic capacity: A systematic review and meta-analysis. *Sports Med* 44(2): 269–279, 2014.
20. González-Alonso J. Hyperthermia impairs brain, heart and muscle function in exercising humans. *Sports Med* 37(4–5): 371–373, 2007.
21. González-Alonso J, Crandall CG, Johnson JM. The cardiovascular challenge of exercising in the heat. *J Physiol* 586(1): 45–53, 2008.
22. Green H, Goreham C, Ouyang J, Ball-Burnett M, Ranney D. Regulation of fiber size, oxidative potential, and capillarization in human muscle by resistance exercise. *Am J Physiol Regul Integr Comp Physiol* 276(2): R591–R596, 1999.
23. Guy JH, Deakin GB, Edwards AM, Miller CM, Pyne DB. Adaptation to hot environmental conditions: an exploration of the performance basis, procedures and future directions to optimise opportunities for elite athletes. *Sports Med* 45(3): 303–311, 2015.
24. Hartmann H, Wirth K, Klusemann M. Analysis of the load on the knee joint and vertebral column with changes in squatting depth and weight load. *Sports Med* 43(10): 993–1008, 2013.
25. Hurst P, Board L. Reproducibility of outdoor 5 km running time-trial in a competitive environment. *Br J Sports Med* 47(17): e4–e4, 2013.
26. Ichinose M, Ichinose-Kuwahara T, Kondo N, Nishiyasu T. Increasing blood flow to exercising muscle attenuates systemic cardiovascular responses during dynamic exercise in humans. *Am J Physiol Regul Integr Comp Physiol* 309(10): R1234–R1242, 2015.
27. Jung AP. The impact of resistance training on distance running performance. *Sports Med* 33(7): 539–552, 2003.
28. Kraemer WJ, Ratamess NA. Fundamentals of resistance training: progression and exercise prescription. *Med Sci Sports Exerc* 36(4): 674–688, 2004.
29. Little JP, Safdar A, Bishop D, Tarnopolsky MA, Gibala MJ. An acute bout of high-intensity interval training increases the nuclear abundance of PGC-1 α and activates mitochondrial biogenesis in human skeletal muscle. *Am J Physiol* 300(6): R1303–R1310, 2011.
30. Ma JK, Scribbans TD, Edgett BA, Boyd JC, Simpson CA, Little JP. Extremely low-volume, high-intensity interval training improves exercise capacity and increases mitochondrial protein content in human skeletal muscle. *Open J Mol Integr Physiol* 2013, 2013.

31. Maughan RJ. Distance running in hot environments: a thermal challenge to the elite runner. *Scand J Med Sci Sports* 20: 95–102, 2010.
32. Maw GJ, Boutcher SH, Taylor NAS. Ratings of perceived exertion and affect in hot and cool environments. *Europ J Appl Physiol* 67(2): 174–179, 1993.
33. McGinley C, Bishop DJ. Influence of training intensity on adaptations in acid/base transport proteins, muscle buffer capacity, and repeated-sprint ability in active men. *J Appl Physiol* 121(6): 1290–1305, 2016.
34. Medicine American College of Sports. *ACSM's Guidelines for Exercise Testing and Prescription*. Lippincott Williams & Wilkins, 2013.
35. Navalta JW, Stone WJ, Lyons ST. Ethical Issues Relating to Scientific Discovery in Exercise Science. *Int J Exerc Sci* 12(1): 1–8, 2019.
36. Neitzel JA, Davies GJ. The benefits and controversy of the parallel squat in strength training and rehabilitation. *Strength Cond J* 22(3): 30, 2000.
37. Nybo L, Nielsen B. Hyperthermia and central fatigue during prolonged exercise in humans. *J Appl Physiol* 91(3): 1055–1060, 2001.
38. Rowell LB, Marx HJ, Bruce RA, Conn RD, Kusumi F. Reductions in cardiac output, central blood volume, and stroke volume with thermal stress in normal men during exercise. *J Clin Invest* 45(11): 1801–1816, 1966.
39. Sjödín B, Jacobs I, Svedenhag J. Changes in onset of blood lactate accumulation (OBLA) and muscle enzymes after training at OBLA. *Eur J Appl Physiol Occup Physiol* 49(1): 45–57, 1982.
40. Sloth M, Sloth D, Overgaard K, Dalgas U. Effects of sprint interval training on VO₂max and aerobic exercise performance: A systematic review and meta-analysis. *Scand J Med Sci Sports* 23(6): e341–352, 2013.
41. Staron RS, Malicky ES, Leonardi MJ, Falkel JE, Hagerman FC, Dudley GA. Muscle hypertrophy and fast fiber type conversions in heavy resistance-trained women. *Eur J Appl Physiol Occup Physiol* 60(1): 71–79, 1990.
42. Tanaka H, Monahan KD, Seals DR. Age-predicted maximal heart rate revisited. *J Am Coll Cardiol* 37(1): 153–156, 2001.
43. Terada S, Kawanaka K, Goto M, Shimokawa T, Tabata I. Effects of high-intensity intermittent swimming on PGC-1 α protein expression in rat skeletal muscle. *Acta Physiol Scand* 184(1): 59–65, 2005.
44. Tucker R, Marle T, Lambert EV, Noakes TD. The rate of heat storage mediates an anticipatory reduction in exercise intensity during cycling at a fixed rating of perceived exertion. *J Appl Physiol* 574(3): 905–915, 2006.
45. Van Der Worp MP, ten Haaf DSM, van Cingel R, de Wijer A, Nijhuis-van der Sanden MWG, Staal JB. Injuries in Runners; A Systematic Review on Risk Factors and Sex Differences. *PloS one* 10(2): e0114937, 2015.
45. Van Gent BRN, Siem DD, van Middelkoop M, van Os TAG, Bierma-Zeinstra SSMA, Koes BBW. Incidence and determinants of lower extremity running injuries in long distance runners: A systematic review. *Br J Sports Med* 41(8): 469–480, 2007.

46. Vanos JK, Warland JS, Gillespie TJ, Kenny NA. Review of the physiology of human thermal comfort while exercising in urban landscapes and implications for bioclimatic design. *Int J Biometeorol* 54(4): 319-334, 2010.
48. Weston AR, Myburgh KH, Lindsay FH, Dennis SC, Noakes TD, Hawley JA. Skeletal muscle buffering capacity and endurance performance after high-intensity interval training by well-trained cyclists. *Eur J Appl Physiol Occup Physiol* 75(1): 7-13, 1996.

