

5-1983

# Food Habits & the Utilization of Drift Organisms by Larval Fishes in the Middle Fork of Drake's Creek, Kentucky

Shirley Timbrook  
*Western Kentucky University*

Follow this and additional works at: <https://digitalcommons.wku.edu/theses>



Part of the [Biology Commons](#)

---

## Recommended Citation

Timbrook, Shirley, "Food Habits & the Utilization of Drift Organisms by Larval Fishes in the Middle Fork of Drake's Creek, Kentucky" (1983). *Masters Theses & Specialist Projects*. Paper 2927.  
<https://digitalcommons.wku.edu/theses/2927>

This Thesis is brought to you for free and open access by TopSCHOLAR®. It has been accepted for inclusion in Masters Theses & Specialist Projects by an authorized administrator of TopSCHOLAR®. For more information, please contact [topscholar@wku.edu](mailto:topscholar@wku.edu).

Timbrook,  
Shirley Kaye

1983

FOOD HABITS AND THE UTILIZATION OF DRIFT ORGANISMS BY LARVAL  
FISHES IN THE MIDDLE FORK OF DRAKE'S CREEK, KENTUCKY

A Thesis

Presented to

the Faculty of the Department of Biology

Western Kentucky University

Bowling Green, Kentucky

In Partial Fulfillment

of the Requirements for the Degree

Master of Science

by

Shirley Kaye Timbrook

May 1983

**AUTHORIZATION FOR USE OF THESIS**

Permission is hereby

granted to the Western Kentucky University Library to make, or allow to be made photocopies, microfilm or other copies of this thesis for appropriate research or scholarly purposes.

reserved to the author for the making of any copies of this thesis except for brief sections for research or scholarly purposes.

Signed

Shelley Long

Date

4-01-83

Please place an "X" in the appropriate box.

This form will be filed with the original of the thesis and will control future use of the thesis.

FOOD HABITS AND THE UTILIZATION OF DRIFT ORGANISMS BY LARVAL  
FISHES IN THE MIDDLE FORK OF DRAKE'S CREEK, KENTUCKY

Recommended March 18, 1983  
(Date)

Robert D. Hoyt  
Director of Thesis

Larry W. Gleason

Blaine R. Ferrell

Approved March 31, 1983  
(Date)

Elmer Gray  
Dean of the Graduate College

TABLE OF CONTENTS

	<u>Page</u>
LIST OF FIGURES .....	iv
LIST OF TABLES .....	v
ACKNOWLEDGEMENTS .....	vi
ABSTRACT .....	vii
INTRODUCTION .....	1
STUDY AREA .....	3
METHODS AND MATERIALS .....	7
RESULTS .....	10
DISCUSSION .....	32
APPENDIX .....	38
LITERATURE CITED .....	43

LIST OF FIGURES

	<u>Page</u>
Figure 1. Map of the Drake's Creek Drainage .....	4
Figure 2. Stream study area of the Middle Fork of Drake's Creek, Kentucky showing Stations I and II .....	5
Figure 3. Percentage frequency of occurrence of various food organisms in the digestive tracts of <i>Hypentelium</i> <i>nigricans</i> during three developmental stages .....	16
Figure 4. Percentage frequency of occurrence of various food organisms in the digestive tracts of <i>Notropis</i> <i>rubellus</i> during three developmental stages .....	18
Figure 5. Percentage frequency of occurrence of various food organisms in the digestive tracts of <i>Notropis</i> <i>cornutus</i> during three developmental stages .....	19
Figure 6. Percentage frequency of occurrence of various food organisms in the digestive tracts of <i>Micropterus</i> sp. during two developmental stages .....	20
Figure 7. Electivity indices "E" for the major food organisms consumed by larval <i>Hypentelium nigricans</i> and <i>Micropterus</i> sp. from 29 April through 20 May and from 7 June through 15 July, respectively .....	22
Figure 8. Electivity indices "E" for the major food organisms consumed by larval <i>Notropis rubellus</i> from 10 May through 29 July, 1982 .....	23
Figure 9. Electivity indices "E" for the major food organisms consumed by larval <i>Notropis cornutus</i> from 10 May through 15 July, 1982 .....	24

LIST OF TABLES

	<u>Page</u>
Table 1. Drift organisms observed in the Middle Fork of Drake's Creek, Kentucky from 18 March through 12 August 1982 .....	11
Table 2. Relative prey density (organisms/m <sup>3</sup> ) of the seven major zooplankton taxa collected in the Middle Fork of Drake's Creek, Kentucky from 29 April through 29 July 1982 .....	12
Table 3. Number of digestive tracts examined and the number of empty tracts observed in each developmental stage for <i>Hypentelium nigricans</i> , <i>Micropterus</i> sp., <i>Notropis cornutus</i> , and <i>Notropis rubellus</i> .....	13
Table 4. Taxa of organisms observed in the stomach contents of <i>Hypentelium nigricans</i> , <i>Micropterus</i> sp., <i>Notropis cornutus</i> and <i>Notropis rubellus</i> .....	15
Table 5. Changes in feeding selectivity of different larval stages of <i>Hypentelium nigricans</i> through time in the Middle Fork of Drake's Creek, 1982 .....	26
Table 6. Changes in feeding selectivity of different larval stages of <i>Notropis rubellus</i> through time in the Middle Fork of Drake's Creek, 1982 .....	27
Table 7. Changes in feeding selectivity of different larval stages of <i>Notropis cornutus</i> through time in the Middle Fork of Drake's Creek, 1982 .....	28
Table 8. Changes in feeding selectivity of different larval stages of <i>Micropterus</i> sp. through time in the Middle Fork of Drake's Creek, 1982 .....	30



## ACKNOWLEDGMENTS

I am indebted to my committee for the successful completion of this study and I wish to express my appreciation to Dr. Robert D. Hoyt, chairman, for his constructive criticism, patience, and understanding; Dr. Larry N. Gleason for his helpful comments and assistance in taxa identification; and Dr. Blaine R. Ferrell for his encouragement and advice throughout the study.

I am grateful to Mr. Rodney McCurry for photographing the figures. I also wish to thank the U. S. Fish and Wildlife Service for providing the plankton nets and anchorage blocks for the project and Western Kentucky University and the Department of Biology for use of facilities and equipment.

A special thanks goes to Mr. Keith Floyd for providing fish specimens for the study.

Funding for this study was provided, in part, through a Graduate Student Research Grant from Western Kentucky University.

FOOD HABITS AND THE UTILIZATION OF DRIFT ORGANISMS BY LARVAL  
FISHES IN THE MIDDLE FORK OF DRAKE'S CREEK, KENTUCKY

Shirley Kaye Timbrook                      May, 1983                      48 pages  
Directed by: R. D. Hoyt, L. N. Gleason, and B. R. Ferrell  
Department of Biology                      Western Kentucky University

Food habits of larvae of the northern hog sucker, common shiner, rosyface shiner, and *Micropterus* sp. from the Middle Fork of Drake's Creek, Kentucky were identified from 18 March to 12 August 1982. Eighteen taxa of animals were observed in the stream drift and a total of seventeen taxa, including larval fish, were identified in the stomach analysis. Eggs, rotifers, hydracarina, and diptera represented the greatest component of the drift comprising 34%, 32%, 8%, and 6%, respectively. The major organisms observed in the gut analysis included eggs (fish, rotifer, and copepod species), rotifers (*Euchlanis* sp.), diptera (Chironomidae), annelida (Naididae), copepoda (cyclopoids), and cladocera (*Alona*, *Camptocercus*, and unknown spp.). The northern hog sucker had the most diverse diet ingesting 15 different taxa, whereas the rosyface shiner consumed only 8 taxa. Rosyface shiners selected for rotifers in their diet during their first four weeks of life and dipterans during the last two weeks as larvae (Ivlev's Electivity Index). Common shiners selected for a greater variety of organisms during their first week while selecting only for dipterans during their last week. Northern hog suckers selected annelids, dipterans, and copepods throughout their larval period. *Micropterus* sp. selected rotifers, annelids, cladocerans, and copepods during their first two weeks and annelids, cladocerans, and

copepods during their last four weeks as larvae. Piscivory was observed in *Micropterus* sp. during the third week of life with 12% of the meta-larvae consuming other larvae. With the exception of *Micropterus* sp., larvae ingested progressively more taxa as they developed from pro- to meta-larvae. The greatest percentage of empty stomachs in all species was observed in the pro-larvae and the fewest in the meta-larvae. Likewise, the rosyface shiner had the greatest percentage of empty stomachs of the four major species studied.

Even as larvae, the four species studied demonstrated resource sharing and positive interrelationships in their food habits.

## INTRODUCTION

Feeding behavior and food habits of fishes have long been of interest to fishery investigators. Such studies provide an understanding of the complex interrelationships of cohabiting fish species in their utilization of stream resources. Relatively few reports, however, have dealt with the feeding habits of larval freshwater stream fishes.

It has been suggested that size-selection predation by larval fish occurs within the morphological limits imposed by mouth size (Northcote 1954, Wong and Ward 1972) and that as mouth size increases with larval fish development there is a selection toward increased prey size (Hartman 1958). Mendelson (1975), however, noted that diet composition may be the result of pressures from others at the same trophic level, and Werner (1979) found that there is a partitioning of food between sympatric fishes. Larkin (1956) in his review of various literature on dietary interrelationships among freshwater fishes concluded that they exercise flexibility in choosing their diets as a result of sharing many available resources, including food, with other species.

The objectives of this study were aimed at identifying the composition of the drift, determining the food habits of the northern hog sucker (*Hypentelium nigricans*), the rosyface shiner (*Notropis rubellus*), the common shiner (*Notropis cornutus*), and *Micropterus* sp., describing the role of drift organisms in the early feeding of fishes, and compiling information on how diet was affected by food availability and other species of fishes in the stream. The four species above were chosen for study

because of their interaction in occupying similar habitats as larvae  
(Floyd 1983).

## STUDY AREA

The Middle Fork of Drake's Creek is a spring-fed stream which arises in Sumner County, Tennessee and flows 33.8 km in a northwesterly direction through Simpson, Allen, and Warren County, Kentucky where it converges with the Trammel Fork and the West Fork to form Drake's Creek (Figure 1). The creek courses through mildly karstic topography characteristic of south central Kentucky. The substrate is chiefly bedrock with intermittent riffle areas of gravel and rubble separating long pools of moving water. Following heavy rains the Middle Fork of Drake's Creek, as well as other streams in the drainage, is subject to a rapid increase in flow rate and water depth but returns to a more normal seasonal flow within a few days.

In this study, two stations were designated at the ford 0.8 km south of Drake, Kentucky, 3.6 km upstream from the mouth of the Middle Fork (Figure 2). Station I was located midstream 4.3 m between the northeast bank and a shoal area and consisted of a hard clay-silt substrate. Station II was situated at the ford 230 m downstream from Station I. It was located midstream 6.5 m between the east and west banks and consisted of a bedrock-hard clay substrate.

During the study period stream flow ranged from 79 cm/sec in March to 10.5 cm/sec in July. Dissolved oxygen concentrations averaged 10.9 mg/l in March decreasing an average of 0.95 mg/l each month thereafter to a minimum of 7.1 mg/l in July. Water depths ranged from a minimum of 25.0 cm and 22.5 cm in July to a maximum of 109.0 cm and 105.0 cm in April at Stations I and II, respectively. However, on at least two occasions in April, the

Figure 1. Map of the Drake's Creek Drainage. The asterisk indicates the collection site on the Middle Fork of Drake's Creek.







Figure 2. Stream study area of the Middle Fork of Drake's Creek,  
Kentucky showing Stations I and II.



stream rose substantially above the 109.0 cm mark and actual depths could not be recorded because of the extremely high water level and rapid current flow in the stream.

## METHODS AND MATERIALS

Drift zooplankton and larval fish collections were made twice weekly from 18 March 1982 through 12 August 1982. Both drift collections and fish samples were taken approximately one hour after analytical sunset.

Drift samples were taken with two 1 m long, 15 cm aperture plankton nets having 0.076 mm mesh. The nets were positioned one above the other on a pair of metal rods anchored in a cement base. Drift samples were of five minute duration at each station site. Larval fish were collected with light traps and with a 3 m long, 0.5 m aperture, conical ichthyoplankton net of 0.5 mm mesh. Light traps were placed in specific habitat areas having minimum current including rooted undercut banks, rock outcroppings and drift-log obstructions while the ichthyoplankton net was placed facing into the current. The light traps were set for 40 minutes while the ichthyoplankton net was set for 5 minutes.

Upon collection, larval fish were immediately fixed in 5% formalin. Drift zooplankton was filtered through a number 20 silk bolting cloth (0.076 mm) and diluted to 40 ml with water to which was added 10 ml of Lugol's Solution. Larval fish were identified using keys of May and Gasaway (1967) and Hogue et al. (1976). Developmental phases (modified from Snyder 1976) were defined as pro-larvae - caudal fin rays not yet apparent; meso-larvae - at least one distinct caudal fin ray apparent but pelvic buds not apparent; meta-larvae - pelvic buds or fins apparent but preanal finfold not entirely absorbed. Five 1 ml samples from each of the four 50 ml drift collections taken on each trip were counted and

the organisms identified using keys of Eddy and Hodson (1970), Peterson (1973), and Pennak (1978). The number of drift organisms per cubic meter of water sampled was determined by averaging the number of organisms from both sampling nets and dividing that number by the volume of water sampled. The volume of water sampled by the drift nets was determined with the aid of a digital flowmeter and the formula

$$\frac{\pi \times (\text{net diameter in meters})^2}{4} \times \frac{\text{flowmeter revolutions} \times 51020}{999,999} \times \text{time}$$

Temperature (degrees Celsius) and dissolved oxygen concentration (mg/l) were determined with a YSI oxygen meter.

Larval and juvenile fishes were randomly selected from each of four species including the northern hog sucker, *Micropterus* sp., the common shiner, and the rosyface shiner. Specimens from each of the pro-larval, meso-larval, and meta-larval developmental stages were chosen for gut analysis for each of the four species. The total length of each fish was recorded to the nearest 0.5 mm with a standard ruler and the gut excised and measured. Contents from the entire digestive tract were removed and permanently mounted in a toluene-butyl acetate-nitrocellulose resin. All food organisms were counted under 600x magnification and identified to genus where possible. Computations of mean numbers of organisms per gut were based on stomachs with and without food. Selection of available food in the environment by fish larvae was determined by use of the quantitative index of electivity, "E", described by Ivlev (1961). The electivity index followed the formula

$$E = \frac{r_i - P_i}{r_i + P_i}$$

where  $E$  = electivity (selectivity)

$r_i$  = % composition of plankter  $i$   
in the ration (gut contents)

and  $p_i$  = % composition of plankter  $i$   
in the food complex (plankton)

"E" values for each major food group occurred within the limits of +1 and -1, the former value indicating complete positive selection and the latter complete rejection of a food item.

## RESULTS

Stream Drift - Eighteen taxa of organisms, in addition to algae, were observed in the stream drift (Table 1). Inter- and intra-station comparisons of drift organisms (Students t,  $P > 0.05$ ) showed no significant difference between the two drift net collections at either station or between stations (Appendix I). Drift organisms were represented primarily by seven groups: eggs (34%), rotifers (32%), hydracarina (8%), dipteran larvae (6%), annelids (2%), copepods (2%), and cladocerans (1%). During the study period, 29 April through 29 July, eggs averaged  $250/m^3$  and constituted 20% to 57% of the total drift component (Table 2). Rotifers exhibited two peaks of abundance, 29 April to 6 May and 21 June to 1 July, during which they averaged  $1065 \text{ organisms}/m^3$ , or 69% of the total. By 29 July, rotifers had decreased to only 3% of the total. Hydracarina increased from 1% on 29 April to 10% on 3 June, reached a maximum of 28% from 7 June to 17 June, then declined to 12% on 29 July. Diptera showed no marked change in percent composition (1%) throughout the study period. Annelids ranged from 0.5% to 5%. Copepods represented no more than 8% of the total throughout the study while cladocerans ranged from  $22/m^3$  (3%) on 1 June to a high of  $111/m^3$  (12%) from 5 July to 15 July.

Stomach Analyses - The number of stomachs examined among the four species included 226 northern hog suckers, 151 *Micropterus* sp., 103 common shiners, and 107 rosyface shiners (Table 3). Rosyface shiners and common shiners exhibited the greatest percentages of empty stomachs with 54% and 42%,

TABLE 1. Drift organisms observed in the Middle Fork of Drake's Creek, Kentucky from 18 March through 12 August 1982.

---



---

EGGS:	CLADOCERA:
Pisces	Alona
Rotatoria	Bosmina longirostris
Copepoda	Bosmina spp.
Unidentified	Camptocercus
GASTROTRICHA	Ceriodaphnia
ROTATORIA:	Daphnia galeata
Asplanchna	Daphnia longispina
Brachionus angularis	Daphnia pulex
Brachionus calyciflorus	Diaphanosoma
Brachionus furculatus	Unidentified spp.
Brachionus havanaensis	NAUPLII
Cephalodella	COPEPODA:
Euchlanis	Calanoida
Filina longiseta	Cyclopoida
Kellicottia longispina	Harpacticoida:
Keratella cochlearis	Canthocamptus
Keratella quadrata	Unidentified
Keratella sp.	HYDRACARINA
Lecane	Hydrachnidae
Lepadella	Unidentified
Monostyla	COLLEMBOLA
Mytilina	PLECOPTERA:
Notholca	Chloroperlidae
Philodina	Allocapnia
Platyias patulus	Perlidae
Polyarthra	Unidentified
Pompholyx	EPHEMEROPTERA
Trichotria	HEMIPTERA
Unidentified spp.	TRICHOPTERA
NEMATODA	COLEOPTERA:
NEMATOMORPHA	Hydrophilidae
TARDIGRADA	Unidentified
ANNELIDA:	DIPTERA:
Oligochaeta:	Chironomidae
Naididae	Heleidae
Unidentified	Simuliidae
ALGAE	Tabanidae

---



TABLE 2. Relative prey density (organisms/m<sup>3</sup>) of the seven major plankton groups collected in the Middle Fork of Drake's Creek, Kentucky from 29 April through 29 July 1982.

T A X A	Organisms/m <sup>3</sup>									
	29 Apr-6 May	10 May-20 May	31 May-3 Jun	7 Jun-17 Jun	21 Jun-1 Jul	5 Jul-15 Jul	29 Jul			
EGGS	308	367	376	130	331	435	138			
ROTATORIA	1062	423	96	70	1068	151	14			
ANNELIDA	25	33	14	6	14	5	24			
CLADOCERA	57	53	22	30	68	111	52			
COPEPODA	7	24	23	4	4	34	36			
HYDRACARINA	17	19	68	126	32	26	53			
DIPTERA	1	19	9	0	1	23	10			

TABLE 3. Number of digestive tracts examined and the number of empty tracts observed in each developmental stage for *Hypentelium nigricans*, *Micropterus* sp., *Notropis cornutus*, and *Notropis rubellus*.

S P E C I E S	EXCISED TRACTS			EMPTY TRACTS				
	Total	Pro	Meso	Meta	Total	Pro	Meso	Meta
<i>Hypentelium nigricans</i>	226	36	116	74	42 (19%)	25 (69%)	17 (15%)	0 (0%)
<i>Micropterus</i> sp.	151	0	75	76	4 (3%)	*	4 (5%)	0 (0%)
<i>Notropis cornutus</i>	103	34	35	34	43 (42%)	33 (97%)	8 (23%)	2 (6%)
<i>Notropis rubellus</i>	107	43	34	30	58 (54%)	30 (70%)	19 (56%)	9 (30%)
Totals	587	113	260	214	147 (25%)	88 (78%)	48 (18%)	11 (5%)

\* No pro-larvae were collected for *Micropterus* sp.

respectively, while the northern hog suckers and *Micropterus* sp. had only 19% and 3%, respectively. The greatest percentage of empty stomachs in all species was observed in the pro-larval and the fewest in the meta-larval stages. In the stomach analysis, a total of 17 taxa, including larval fish and algae, were identified among the four larval species. The major prey groups observed included eggs (fish, copepod, and rotifer species), rotifers (*Euchlanis* sp.), diptera (Chironomidae), annelids (Naididae), copepods (cyclopoids), and cladocerans (*Alona*, *Camptocercus*, and unknown species) (Table 4). The northern hog sucker had the most diverse diet ingesting 15 different taxa (Tables 1 & 4). *Micropterus* sp. (Tables 1 & 4) and the common shiner (Tables 1 & 4) consumed 13 and 11 taxa, respectively, while the rosyface shiner (Tables 1 & 4) ingested 8 taxa. However, rosyface shiners contained prominent algal masses in their stomach contents.


Pro-larvae - Most pro-larval representatives of the four species studied (78%) contained yolk material in the digestive tract with no food organisms (Table 3). However, some pro-larval specimens of each species, with the exception of *Micropterus* sp. which had no pro-larval representatives, contained food organisms in addition to yolk. The maximum total lengths of pro-larvae containing yolk were 14.75 mm, 6.50 mm, and 6.00 mm for the northern hog sucker, the rosyface shiner, and the common shiner, respectively. Thirty-one percent of pro-larval northern hog suckers ingested four groups of food organisms including eggs, rotifers, cladocerans, and hydracarina (Figure 3). Rosyface shiner pro-larvae

TABLE 4. Food items and taxa of organisms observed in the stomach contents of *Hypentelium nigricans*, *Micropterus* sp., *Notropis cornutus*, and *Notropis rubellus*.

FOOD/PREY T A X A	F I S H T A X A			
	<i>Hypentelium nigricans</i>	<i>Micropterus sp.</i>	<i>Notropis cornutus</i>	<i>Notropis rubellus</i>
EGGS	+	+	+	-
GASTROTRICHA	-	-	+	-
ROTATORIA	+	+	+	+
NEMATODA	+	-	+	-
ANNELIDA	+	+	+	+
ALGAE	+	+	+	+
CLADOCERA	+	+	+	+
NAUPLII	+	+	+	-
COPEPODA	+	+	+	+
HYDRACARINA	+	+	+	+
COLLEMBOLA	+	-	-	-
PLECOPTERA	+	+	-	-
EPHEMEROPTERA	+	-	-	+
COLEOPTERA	+	+	-	-
DIPTERA	+	+	+	+
NEUROPTERA	+	+	-	-
PISCES	-	+	-	-

Figure 3. Percentage frequency of occurrence of various food organisms in the digestive tracts of *Hypentelium nigricans* during three developmental stages.

Hypentelium nigricans



ingested rotifers (*Euchlanis* sp.) and dipteran larvae (Chironomidae) (Figure 4) while common shiner pro-larvae (Figure 5) ingested only rotifers (*Euchlanis* sp.). Most of the pro-larvae of all species containing stomach contents also included various algal forms (Table 4).

Meso-larvae - Meso-larval stages in all species had some representatives which still contained yolk material in their guts. The maximum total lengths of meso-larvae containing yolk were 14.25 mm, 8.00 mm, 7.00 mm, and 8.50 mm for the northern hog sucker, the rosyface shiner, the common shiner, and *Micropterus* sp., respectively. Meso-larvae of northern hog suckers and common shiners included representatives of all seven major prey groups in their stomach contents (Figures 3 & 5). The four most abundant groups in both species included rotifers, dipteran larvae, eggs, and annelids. *Micropterus* sp. ingested six major prey items with very high percent frequencies of occurrence for cladocerans, copepods, and dipteran larvae but with no hydracarina (Figure 6). Rosyface shiner meso-larvae contained representatives of only three taxa: rotifers, dipterans, and hydracarina (Hydrachnid) (Figure 4).

Meta-larvae - Meta-larval representatives of all four species, except the rosyface shiner, ingested specimens of all seven major food categories (Figures 3 - 6). Dipteran larvae represented the greatest frequency of occurrence in all species observed. Northern hog suckers showed the greatest balance in stomach contents with over 40% of the stomachs containing organisms of every food group except hydracarina (Figure 3). *Micropterus* sp. showed very high percentage frequencies for dipterans and copepods (Figure 6). Rosyface shiners indicated the least complete diet

Figure 4. Percentage frequency of occurrence of various food organisms in the digestive tracts of *Notropis rubellus* during three developmental stages.






Figure 5. Percentage frequency of occurrence of various food organisms in the digestive tracts of *Notropis cornutus* during three developmental stages.





Figure 6. Percentage frequency of occurrence of various food organisms in the digestive tracts of *Micropterus* sp. during two developmental stages.


Micropterus sp.

Meso-larvae N = 75



Micropterus sp.

Meta-larvae N = 76



of all the four species with less than 10% frequency of occurrence for all but one dietary component, dipteran larvae (Figure 4).


With the exception of *Micropterus* sp., larvae ingested progressively more food groups as they developed from pro- to meta-larval stages (Appendices II - V). Piscivory was observed in *Micropterus* sp. during the third week of life with 12% of the meta-larvae consuming other fish larvae. The identifiable larval fish ingested were *Notropis* spp., and those which could be specifically identified were *N. rubellus*. It was observed that at this developmental stage, *Micropterus* sp., as opposed to the other non-piscivorous species, had a highly coiled digestive tract which allowed for accomodation of fish larvae equal to or exceeding its own body length (11.5 mm TL at the onset of piscivory). The shiner's digestive tract remained straight and the northern hog sucker, though it had a straight gut, had a greater gut-to-total-length ration (Appendices II - V).

Generally, size of prey organisms increased from pro-larval to meta-larval stages. However, in each developmental stage, the prey taxa consumed were usually the same, i.e. *Euchlanis* sp. was the major rotifer observed in all species in all developmental stages. As fish species increased in size, dipterans increased in size from exclusively larval forms in meso-larval stomachs to larvae and pupae in meta-larval guts.

Electivity Indices - Considerable variation was observed among the four fish species as to their selection for and against the seven major food groups throughout the larval period. Indices of electivity (Figures 7 - 9) showed that northern hog suckers selected for annelids, copepods, and

Figure 7. Electivity indices, "E", for the major food organisms consumed by larval *Hypentelium nigricans* and *Micropterus* sp. from 29 April through 20 May and from 7 June through 15 July, respectively.

*Hypentelium nigricans* N = 226



*Micropterus* sp. N = 151






Figure 8. Electivity indices, "E", for the major food organisms consumed by larval *Notropis rubellus* from 10 May through 29 July 1982.

Notropis rubellus

N=107





Figure 9. Electivity indices, "E", for the major food organisms consumed by larval *Notropis cornutus* from 10 May through 15 July 1982.

Notropis cornutus

N=103



dipterans and against cladocerans and rotifers, despite their high frequency of occurrence in the diet (Figure 7). Rosyface shiners selected for rotifers in their diet during their first four weeks of life, increasing their selectivity to annelids, cladocerans, copepods, and dipterans during the seventh and eighth week and finally selecting only for dipterans during their last week as larvae (Figure 8). The common shiner selected for a wide variety of organisms during the first six weeks, including rotifers, annelids, and dipterans, while selecting only for dipterans during the last week as larvae (Figure 9). Basically, *Micropterus* sp. selected for annelids, cladocerans, and copepods throughout the larval stages (Figure 7). Comparing dates of prey selection, *Micropterus* sp. decreased the selectivity for cladocerans and dipterans on 5 July (Figure 7) when the rosyface shiner began selecting these taxa (Figure 8).

Pro-larvae - As pro-larvae, the northern hog sucker selected against the major food groups during the first two weeks and selected for copepods, hydracarina, and cladocerans during the second two weeks (Table 5). No pro-larvae were available for analysis of *Micropterus* sp. Both the rosyface shiner (Table 6) and the common shiner (Table 7) selected rotifers during their first two weeks as pro-larvae. In addition, the rosyface shiner selected dipteran larvae which became the only food item selected during the second two weeks. From the sixth week through the end of the pro-larval period, the rosyface shiner selected against all seven food categories. The general pro-larval selectivity pattern, therefore, showed

TABLE 5. Changes in feeding selectivity of different larval stages of *Hypentelium nigricans* through time in the Middle Fork of Drake's Creek, 1982.

LIFE STAGE	PREY	Apr 29 - May 6			May 10 - May 20			Electivity Index (E)
		% in gut (r)	% in nets (p)	Electivity Index (E)	% in gut (r)	% in nets (p)	Electivity Index (E)	
<b>Pro-larvae:</b>								
	EGGS	0	20.00	-1.00	21.05	35.00	-0.24	
	ROTATORIA	0	69.00	-1.00	10.52	40.00	-0.58	
	ANNELIDA	0	2.00	-1.00	0	3.00	-1.00	
	COPEPODA	0	0.49	-1.00	10.52	2.00	+0.68	
	CLADOCERA	0	4.00	-1.00	5.26	5.00	+0.25	
	HYDRACARINA	0	1.00	-1.00	5.26	2.00	+0.44	
	DIPTERA	0	0.04	-1.00	0	2.00	-1.00	
<b>Meso-larvae:</b>								
	EGGS	14.62	20.00	-0.15	13.31	55.00	-0.44	
	ROTATORIA	17.54	69.00	-0.59	11.41	40.00	-0.55	
	ANNELIDA	25.73	2.00	+0.85	12.17	3.00	+0.60	
	COPEPODA	3.51	0.49	+0.75	5.32	2.00	+0.45	
	CLADOCERA	0	4.00	-1.00	0.95	5.00	-0.68	
	HYDRACARINA	1.17	1.00	+0.07	1.71	2.00	-0.07	
	DIPTERA	16.96	0.04	+0.99	28.33	2.00	+0.86	
<b>Meta-larvae:</b>								
	EGGS	9.68	20.00	-0.34	10.31	35.00	-0.54	
	ROTATORIA	19.35	69.00	-0.56	16.41	40.00	-0.41	
	ANNELIDA	22.58	2.00	+0.83	10.55	3.00	+0.55	
	COPEPODA	3.22	0.49	+0.73	5.23	2.00	+0.44	
	CLADOCERA	0	4.00	-1.00	1.84	5.00	-0.46	
	HYDRACARINA	0	1.00	-1.00	1.26	2.00	-0.22	
	DIPTERA	3.22	0.04	+0.97	28.27	2.00	+0.86	

Table 6. Changes in feeding selectivity of different larval stages of *Notropis rubellus* through time in the Middle Fork of Drakes' Creek, 1982.

TAXON	May 10 - May 20			Jun 7 - Jun 17			Jun 21 - Jul 1			Jul 5 - Jul 15			Jul 29		
	%/gut (r)	%/net (p)	Electiv. Index (E)	%/gut (r)	%/net (p)	Electiv. Index (E)	%/gut (r)	%/net (p)	Electiv. Index (E)	%/gut (r)	%/net (p)	Electiv. Index (E)	%/gut (r)	%/net (p)	Electiv. Index (E)
Pro-larvae:															
EGGS	0	55.00	-1.00	0	28.00	-1.00	0	21.00	-1.00	0	49.00	-1.00	0		
ROTATORIA	91.00	40.00	+0.40	63.16	15.00	+0.61	61.11	69.00	-0.06	0	17.00	-1.00	0		
ANNELIDA	0	5.00	-1.00	0	1.00	-1.00	0	1.00	-1.00	0	0.56	-1.00	0		
COPEPODA	0	2.00	-1.00	0	0.88	-1.00	0	0.50	-1.00	0	4.00	-1.00	0		
CLADOCERA	0	5.00	-1.00	0	6.00	-1.00	0	4.00	-1.00	0	12.00	-1.00	0		
HYDRACARINA	0	2.00	-1.00	0	28.00	-1.00	0	2.00	-1.00	0	3.00	-1.00	0		
DIPTERA	5.88	2.00	+0.49	0	0.05	-1.00	0	0.05	-1.00	0	2.00	-1.00	0		
no pro-larvae taken															
Meso-larvae:															
EGGS	0	55.00	-1.00	0	28.00	-1.00	0	21.00	-1.00	0	49.00	-1.00	0		
ROTATORIA	0	40.00	-1.00	65.45	15.00	+0.62	68.75	69.00	0	0	17.00	-1.00	0		
ANNELIDA	0	5.00	-1.00	0	1.00	-1.00	0	1.00	-1.00	0	0.56	-1.00	0		
COPEPODA	0	2.00	-1.00	0	0.88	-1.00	0	0.50	-1.00	0	4.00	-1.00	0		
CLADOCERA	0	5.00	-1.00	0	6.00	-1.00	0	4.00	-1.00	0	12.00	-1.00	0		
HYDRACARINA	0	2.00	-1.00	0	28.00	-1.00	0	2.00	-1.00	0	3.00	-1.00	0		
DIPTERA	0	2.00	-1.00	0	0.05	-1.00	0	0.05	-1.00	0	2.00	+0.96	100.00	0	
no meso-larvae taken															
Meta-larvae:															
EGGS	0	55.00	-1.00	0	28.00	-1.00	0	21.00	-1.00	0	49.00	-1.00	0		
ROTATORIA	0	40.00	-1.00	65.45	15.00	+0.62	68.75	69.00	0	0	17.00	-1.00	0		
ANNELIDA	0	5.00	-1.00	0	1.00	-1.00	0	1.00	-1.00	0	0.56	-1.00	0		
COPEPODA	0	2.00	-1.00	0	0.88	-1.00	0	0.50	-1.00	0	4.00	-1.00	0		
CLADOCERA	0	5.00	-1.00	0	6.00	-1.00	0	4.00	-1.00	0	12.00	-1.00	0		
HYDRACARINA	0	2.00	-1.00	0	28.00	-1.00	0	2.00	-1.00	0	3.00	-1.00	0		
DIPTERA	0	2.00	-1.00	0	0.05	-1.00	0	0.05	-1.00	0	2.00	+0.92	100.00	0	
no meta-larvae taken															
EGGS	0	55.00	-1.00	0	28.00	-1.00	0	21.00	-1.00	0	49.00	-1.00	0		
ROTATORIA	0	40.00	-1.00	65.45	15.00	+0.62	68.75	69.00	0	0	17.00	-1.00	0		
ANNELIDA	0	5.00	-1.00	0	1.00	-1.00	0	1.00	-1.00	0	0.56	-1.00	0		
COPEPODA	0	2.00	-1.00	0	0.88	-1.00	0	0.50	-1.00	0	4.00	-1.00	0		
CLADOCERA	0	5.00	-1.00	0	6.00	-1.00	0	4.00	-1.00	0	12.00	-1.00	0		
HYDRACARINA	0	2.00	-1.00	0	28.00	-1.00	0	2.00	-1.00	0	3.00	-1.00	0		
DIPTERA	0	2.00	-1.00	0	0.05	-1.00	0	0.05	-1.00	0	2.00	+0.92	45.45	0	





no overlapping among the four species. The only exception was the selection for rotifers by both the rosyface shiner and the common shiner. However, selection for this taxon occurred during a time when rotifers constituted 40% of the community (Table 2).

Meso-larvae - Meso-larval northern hog suckers (Table 5) selected for dipterans, annelids, and copepods throughout this four week developmental stage eliminating hydracarina as a preferred food item after the first two weeks. The rosyface shiner meso-larvae responded negatively to most food items with a preference for rotifers by the fifth week and a selection for dipterans by the sixth week (Table 6) while common shiner meso-larvae selected for dipterans, annelids, and rotifers (Table 7). *Micropterus* sp. selected for dipterans, copepods, annelids, and rotifers as young meso-larvae and for copepods and dipterans as late meso-larvae (Table 8). Though meso-larval northern hog suckers (Table 5) continued to select annelids, that selection decreased on 10 May when the common shiner meso-larvae (Table 7) started feeding on the same food item.

Meta-larvae - Meta-larval northern hog suckers continued to select for copepods in their diet as they did throughout earlier stages (Table 5). In addition, dipterans and annelids were also selected. Rosyface shiner meta-larvae had a high selectivity for dipterans with only annelids and copepods appearing in the diet at mid-meta-larval stages (Table 6). Common shiner meta-larval food preferences changed throughout this developmental stanza. While dipteran larvae were selected for throughout the period, alternate selections for cladocerans, annelids, rotifers, copepods, miscellaneous eggs, and hydracarina were observed (Table 7).

TABLE 8. Changes in feeding selectivity of different larval stages of *Micropterus* sp. through time in the Middle Fork of Drake's Creek, 1982.

LIFE STAGE	PREY	Jun 7 - Jun 17		Jun 21 - Jul 1		Jul 5 - Jul 15				
		% in gut (r)	% in nets (p)	Electiv. Index (E)	% in gut (r)	% in nets (p)	Electiv. Index (E)	% in gut (r)	% in nets (p)	Electiv. Index (E)
Pro-larvae:		no pro-larvae taken		no pro-larvae taken		no pro-larvae taken		no pro-larvae taken		
Meso-larvae:		no pro-larvae taken		no pro-larvae taken		no pro-larvae taken		no meso-larvae taken		
EGGS		7.34	28.00	-0.58	6.10	21.00	-0.54			
ROTATORIA		20.21	15.00	+0.14	0	69.00	-1.00			
ANNELIDA		4.57	1.00	+0.64	0	1.00	-1.00			
COPEPODA		33.81	0.88	+0.94	82.44	0.29	+0.99			
CLADOCERA		19.61	6.00	+0.53	1.52	4.00	-0.44			
HYDRACARINA		0	28.00	-1.00	0	2.00	-1.00			
DIPTERA		9.02	0.05	+0.98	6.87	0.05	+0.98			
Meta-larvae:		no meta-larvae taken		no meta-larvae taken		no meta-larvae taken		no meta-larvae taken		
EGGS										
ROTATORIA					7.00	21.00	-0.50	3.22	49.00	-0.87
ANNELIDA					1.83	69.00	-0.94	0	17.00	-1.00
COPEPODA					1.29	1.00	+0.12	9.67	0.56	+0.89
CLADOCERA					49.01	0.29	+0.98	38.70	4.00	+0.81
HYDRACARINA					10.60	4.00	+0.45	16.13	12.00	+0.14
DIPTERA					0.33	2.00	-0.71	3.22	3.00	+0.03
					20.87	0.05	+0.99	16.13	2.00	+0.77

*Micropterus* sp. had the greatest selection pattern of the four species by positively selecting for four taxa as early meta-larvae and five taxa during later meta-larval development (Table 8).

Generally, the four fish species selected for more and larger prey organisms as they advanced from pro-larval to meta-larval stages.

## DISCUSSION

Invertebrate drift is a normal feature of lotic systems (Elliott and Corlett 1971). However, Hynes (1970) found that these type environments are poorly suited for the production and maintenance of zooplankton and that streams consequently harbor small zooplankton communities, especially microcrustaceans. His observations were similar to those of this study wherein copepods and cladocerans comprised the smallest percentage of organisms in the community. Rotifers, on the other hand, constituted a high percentage which supported the findings of Winner (1975) that these organisms are found in abundance in lotic environments and often dominate the zooplankton community. It does not appear unusual that eggs would also occupy a high percentage level since they constitute an unresisting status in a moving water body and are consequently swept into the drift. Seasonal abundance peaks are common, if not universal, in the zooplankton community and several studies have reported the occurrence (Pennak 1949, Hudson and Cowell 1966, Elliott and Corlett 1971, and Selgeby 1975). The present study was no exception, for rotifers exhibited two distinct peaks in abundance during the study period.

The observation that pro-larval fishes in Drake's Creek had a higher percentage of empty digestive tracts than fishes of a more advanced developmental stage was similar to that reported by Siefert (1972). This find was suggested by Siefert (1972) to be a result of incompletely developed digestive tracts in smaller larvae. Of the four major species studied from Drake's Creek, rosyface shiners and common shiners exhibited

the greatest percentage of empty digestive tracts. Both species exhibited intestines that remained long and straight with little observable morphometric differentiation throughout the larval period. On the other hand, northern hog suckers and *Micropterus* sp. had low percentages of empty guts. The northern hog sucker had a greater gut-to-total length ratio than did the shiners. *Micropterus* sp. demonstrated a highly coiled digestive tract with the loops dividing the gut into definite functional parts.

The utilization of members of 17 different taxa of food organisms by the four species from Drake's Creek was a surprising observation. While it is well known that many adult fishes feed upon a wide variety of organism types, selecting at particular times for individual taxa, the extension of that premise to larval feeding has not been verified and, consequently, was surprising. The greatest feeding diversity observed by the northern hog sucker in Drake's Creek was similar to the findings reported by Siefert (1972) for the white sucker (*Catostomus commersoni*). Pflieger (1975) also reported the northern hog sucker to be an aggressive feeder.

Yolk material was found to occur in pro-larval and meso-larval digestive tracts, often in addition to extraneous food organisms. Lagler et al. (1977) found that many young fishes with tiny mouths began to feed on plankton while still containing yolk bodies. In their study on larval drum (*Aplodinotus grunniens*), Clark and Pearson (1979) observed feeding to commence well before the yolk sac was absorbed and that 26% of the specimens examined had both food and yolk material in the gut. Furthermore,

the combination of extraneous food and yolk in the digestive tract has been reported in the spotted sucker (*Minytrema melanops*) (White and Haag 1977) and the walleye (*Stizostedion vitreum*) (Bulkley et al. 1976). As pointed out by Clark and Pearson (1979), it is not unusual for the larval fish to ingest foods before entire absorption of the yolk considering that mouthparts and an oral opening form during the pro-larval stage when the yolk sac is still being absorbed.

As fish size increases, the size, variety, and number of food organisms generally increases. That smaller larvae utilize smaller prey and larger larvae utilize larger prey was illustrated in this study by smaller pro-larvae consuming rotifers while meso- and meta-larvae consumed dipteran larvae. Similar findings have been reported by Silvellis (1949), Ivlev (1961), Hickling (1966), Siefert (1968), Scalet (1972), Mathur (1977), Werner (1979), Overmann et al. (1980), Applegate (1981), and Watkins et al. (1981). This is a general phenomenon for many fishes (Hynes 1970). Studies have suggested that mouth sizes determine the size of prey taken (Northcote 1954, Wong and Ward 1972) and that, within a species, as mouth size increases prey size increases (Hartman 1958).

The use of a greater variety of food organisms by larvae as they increased in size in Drake's Creek was similar to the findings of Heins and Clemmer (1975) and Overmann et al. (1980). Wallace (1976) proposed that the use of a larger variety of food with increased fish size could be explained by the increased range of food sizes available with increased mouth size rather than by a change in feeding behavior. Another reason

for the shift in preferred food items might be, as reported by Nilsson (1965), that as fish size increases, energy expended in capturing smaller prey becomes uneconomical. This alteration in food habits of larval fishes becomes important as size dependent differences in feeding reduce interspecific competition (Hynes 1970) and allows for more efficient utilization of the habitat resources. Other studies have also shown that fish frequently select the largest species of zooplankton and the largest individuals of those species (Brooks and Dodson 1965, Galbraith 1967). Learned behavior and previous experience have been suggested as reasons for an individual to continue to choose an item in the diet (Ware 1971, Wong and Ward 1972, Hansen and Wahl 1981).

Twelve percent of the *Micropterus* sp. meta-larvae demonstrated piscivory. A review of the literature provided few reports of piscivory with fish less than 20 mm total length. Clark and Pearson (1979) reported 3.8 mm freshwater drum to consume other larvae. Piscivory was reported for 8.0 mm walleye (Bulkley et al. 1976) and for 17 mm longnose gar (*Lepososteus osseus*) (Echelle 1968). Therefore, the Drake's Creek finding of 11.5 mm larvae becoming piscivorous is one of few such observations. There have been reports of cannibalism by *Micropterus* sp. at 18 mm (Chew 1974) and at 23.6 mm (Krammer and Smith (1960) but no cannibalism was found in the Drake's Creek study. The great abundance of shiners in the stream (Floyd 1983) probably explained their utilization by the *Micropterus* sp.

Selection for or against a particular food organism has been reported

for a wide variety of fishes. Food habits found in this study suggested that a degree of feeding specialization existed among the larval fish studied. Pro-larvae of each species selecting for different organisms than pro-larvae of the other species supported this contention. This observation of different diet selection by like larval stages of different species was also seen by Bulkley et al. (1976) in their study of cohabiting fishes. Likewise, the Drake's Creek finding of the species, rather than the developmental stage, changing its selection of a food item when another species began selecting the same item suggested a resource sharing adjustment. Bulkley et al. (1976) reported the same observation with other cohabiting species. The Drake's Creek findings of feeding specialization by larval members of different species seems to support resource partitioning as reported by Wynes and Wissing (1982) and George and Hadley (1979), the competitive exclusion principal (Elton 1946, Zaret and Rand 1971), and the existence of positive interrelationships among cohabiting species (Brocksen et al. 1968, Mendelson 1975).

Many authors have postulated as to why fishes select for or against certain food organisms. Brooks and Dodson (1965) suggested size, abundance, and the ability to avoid a predator as properties of a food organism which cause it to be selected for or against. Lagler et al. (1977) also suggested prey abundance as a determining factor in selection. However, the present study showed dipteran larvae to rank low in abundance but yet were selected for by all species during all developmental stages. Siefert (1972) studied shiners and centrarchids and found that early larvae selected food



organisms that were small enough and slow enough to be desirable by that group. In addition, Moore and Moore (1976) found that prey organisms probably attract predators by their movements such as cyclopoid copepods swimming by pronounced jerks. Cyclopoid copepods were among the first food items of shiner and sucker larvae in this study as in the Moore and Moore (op cit) study.

Ivlev (1961) reported experience to be a significant influence on prey selection by fishes. Brooks (1967) observed fish to continue to select small prey even after a larger prey type becomes more abundant. Electivity data in this study also showed a continuation in selecting for the same species in later larval stages. Since attack distance of fish is related to prey size (Confer et al. 1978) and dependent on experience (Ware 1971), fish may be more effective at capturing taxa with which they are familiar (Hansen and Wahl 1981). Wong and Ward (1972) also found that experience may induce a continued utilization of smaller, more familiar prey organisms.

Appendix I. Drift organisms/m<sup>3</sup>, through time, for Station I-A (upper net) and Station I-B (lower net), for Station II-A (upper net) and Station II-B (lower net) and for Stations I and II.

		Sta. I-A	Sta. I-B	Sta. II-A	Sta. II-B	Sta. I	Sta. II
Mar.	18	285.24	389.73	586.96	1279.36	337.48	933.16
	22	305.35	578.56	269.48	759.73	441.95	514.60
	24	398.02	472.65	393.66	681.00	435.33	537.33
	29	98.03	141.17	459.30	337.21	119.60	398.25
Apr.	1	853.44	1030.15	1882.98	1854.61	941.79	1868.79
	5	*	*	383.01	54.49		
	10	2848.26	2914.21	1319.16	2088.36	2881.24	1703.76
	12	981.03	1528.00	1228.30	518.55	1254.51	873.43
	15	2216.90	3066.19	2317.93	2279.53	2641.54	2298.73
	19	538.14	1234.14	*	*		
	26	1409.36	1473.91	797.95	793.29	1441.63	795.62
	29	473.20	445.36	985.91	889.86	459.28	937.89
May	4	2015.60	2274.22	2057.52	2297.64	2144.91	2177.58
	6	1507.66	1932.78	1686.44	1745.77	1720.22	1716.11
	10	1930.25	1484.18	2018.28	1731.06	1707.22	1874.67
	13	490.55	332.87	751.84	516.89	411.71	634.36
	17	957.04	588.23	1238.53	684.23	772.64	961.38
	20	922.49	1118.53	*	*		
	24	710.90	663.75	398.06	432.94	687.32	415.50
	27	601.39	311.19	397.97	305.70	456.29	351.84
	31	1312.26	1717.59	653.64	505.84	1514.92	579.74
	Jun	3	567.81	227.11	437.25	182.37	397.46
7		490.57	219.76	508.14	289.79	355.16	398.97
10		451.67	162.60	420.90	212.88	307.13	316.89
14		346.82	352.59	480.46	228.01	349.70	354.23
17		577.21	307.35	683.53	642.51	442.28	663.02
21		996.92	585.81	796.73	349.96	791.36	573.34
28		2491.34	1328.71	5207.68	2779.52	1910.03	3993.60
Jul.	1	1465.90	897.30	1236.28	368.89	1181.60	802.59
	5	1067.65	221.99	94.23	231.65	644.82	162.94
	8	752.08	1121.17	705.38	255.58	936.63	480.48
	12	2820.66	1600.45	774.62	2499.98	2210.56	1637.30
	15	550.66	539.53	692.19	405.00	545.10	548.60
	19	241.35	482.71	194.83	80.22	362.03	137.53
	22	294.11	741.97	275.37	423.26	518.04	349.31
	26	432.92	189.04	725.21	153.21	310.98	439.21
	29	580.11	572.63	293.38	276.85	576.37	285.11
Aug.	9	517.40	254.00	92.66	85.53	385.70	89.10
	12	92.50	240.50	86.31	43.15	166.50	64.73

\* No data available

Appendix II. Total length category, number of individuals, mean total length, mean gut length, gut length-total length ratio, and mean number of food organisms per stomach contents for each of three developmental stages of *Hypentelium nigricans* from the Middle Fork of Drake's Creek, Kentucky. (E=eggs, R=rotifers, A=annelids, Cop=copepods, Clad=cladocerans, H=hydracarina, and D=dipterans).

Life Stage	Total length category (mm)	N	Mean TL	Mean gut								
				length (mm)	Gut:TL	E	R	A	Cop	Clad	H	D
Pro-larvae	10.00-11.99	18	11.00	5.93	0.54:1	0.05	0.11	0	0	0	0	0
	12.00-13.99	9	13.64	7.36	0.54:1	0	0.22	0	0.11	0	0.11	0
	14.00-15.99	9	14.22	7.22	0.51:1	0	0.11	0	0	0.22	0	0
Meso-larvae	10.00-15.99	36	12.46	6.61	0.53:1	0.02	0.14	0	0.03	0.05	0.03	0
	10.00-11.99	14	11.25	6.21	0.55:1	0.36	0.36	0.36	0.14	0.07	0	0.93
	12.00-13.99	12	12.89	6.54	0.51:1	0.33	0.50	0.25	0.33	0.17	0	1.17
	14.00-15.99	81	14.88	7.91	0.53:1	0.75	0.53	0.52	0.21	0.10	0.11	1.65
Meta-larvae	16.00-17.99	9	16.30	8.00	0.49:1	1.33	1.89	3.55	0.78	0.11	0.22	2.11
	10.00-17.99	116	14.35	7.57	0.53:1	0.70	0.61	0.71	0.26	0.10	0.09	1.55
	15.00-16.99	19	16.08	8.06	0.50:1	3.00	2.21	2.42	1.21	0.26	0.26	6.95
Meta-larvae	17.00-18.99	41	17.78	8.63	0.48:1	3.46	6.07	3.93	1.56	0.46	0.32	7.66
	19.00-21.00	14	19.64	9.32	0.47:1	1.78	5.07	2.43	1.36	0.64	1.43	8.71
Meta-larvae	15.00-21.00	74	17.69	8.61	0.49:1	3.02	4.89	3.26	1.43	0.44	0.51	7.68

Appendix III. Total length category, number of individuals, mean total length, mean gut length, gut length-total length ratio, and mean number of food organisms per stomach contents for each of three developmental stages of *Notropis rubellus* from the Middle Fork of Drake's Creek, Kentucky. (E=eggs, R=rotifers, A=annelids, Cop=copepods, Clad=cladocerans, H=hydracarina, and D=dipterans).

Life Stage	Total length category (mm)	N	Mean TL	Mean gut									
				Gut:TL	E	R	A	Cop	Clad	H	D		
Pro-larvae	5.00- 6.49	22	6.05	2.00	0.33:1	0	0.41	0	0	0	0	0	0
	6.50- 7.99	21	6.76	2.34	0.35:1	0	1.19	0	0	0	0	0	0.05
	5.00- 7.99	43	6.40	2.17	0.34:1	0	0.79	0	0	0	0	0	0.02
Meso-larvae	6.00- 7.49	24	6.77	2.16	0.32:1	0	11.50	0	0	0	0	0	0
	7.50- 9.99	10	8.15	2.40	0.29:1	0	10.50	0	0	0	0	1.00	3.00
	6.00- 9.99	34	7.17	2.23	0.31:1	0	11.20	0	0	0	0	0.29	0.88
Meta-larvae	10.00-11.99	5	10.85	3.70	0.34:1	0	0	0	0	0	0	1.00	1.00
	12.00-13.99	12	12.67	4.43	0.35:1	0	0	1.00	1.00	0	0	1.00	2.00
	14.00-15.99	5	15.30	5.95	0.39:1	0	0	1.00	0	0	0	0	1.00
	16.00-17.99	3	16.58	7.00	0.42:1	0	0	0	0	0	0	0	1.50
	18.00-19.99	3	18.08	8.58	0.47:1	0	0	0	0	1.00	0	0	1.00
	22.00-27.99	2	24.25	12.00	0.49:1	0	0	0	0	0	0	0	0
10.00-27.99	30	14.51	5.74	0.40:1	0	0	0	0.57	0.50	0	0.57	1.38	

Appendix IV. Total length category, number of individuals, mean total length, mean gut length, gut length-total length ratio, and mean number of food organisms per stomach contents for each of three developmental stages of *Notropis cornutus* from the Middle Fork of Drake's Creek, Kentucky. (E=eggs, R=rotifers, A=annelids, Cop=copepods, Clad=cladocerans, H=hydracarina, and D=dipterans).

Life Stage	Total length category (mm)	N	Mean TL (mm)	Mean gut									
				Gut:TL	E	R	A	Cop	Clad	H	D		
Pro-larvae	5.00- 5.99	26	5.45	2.13	0.39:1	0	0	0	0	0	0	0	
	6.00- 7.99	8	6.16	2.69	0.44:1	0	0.12	0	0	0	0	0	
	5.00- 7.99	34	5.62	2.26	0.40:1	0	0.01	0	0	0	0	0	
Meso-larvae	6.00- 7.99	8	7.06	2.93	0.42:1	0	0.37	0	0	0	0	0.25	
	8.00- 9.99	20	9.05	3.73	0.41:1	0.50	11.75	1.10	0.05	0.15	0	0.75	
	10.00-11.99	4	10.31	4.25	0.41:1	0	1.00	2.00	0.25	0.50	0	2.00	
	12.00-13.99	3	12.17	4.17	0.34:1	0	0	0	0	0	0	0.53	2.00
	6.00-13.99	35	9.01	3.64	0.40:1	0.28	6.91	0.86	0.06	0.14	0.03	0.88	
Meta-larvae	9.00-12.99	4	10.94	4.56	0.42:1	0	0.25	1.50	0.50	0.75	0.25	6.25	
	13.00-16.99	13	14.40	6.54	0.45:1	0.08	0.92	0.15	0	0	0.69	4.46	
	17.00-20.99	14	18.39	12.09	0.66:1	0	0.36	0.36	0	0	0.07	2.14	
	21.00-24.99	3	21.17	9.75	0.46:1	0	0.33	0.33	0	0	0	2.00	
	9.00-24.99	34	16.23	8.87	0.55:1	0.03	0.56	0.41	0.06	0.09	0.32	3.50	

Appendix V. Total length category, number of individuals, mean total length, mean gut length, gut length-total length ratio, and mean number of food organisms per stomach contents for each of two developmental stages of *Micropotenus* sp. from the Middle Fork of Drake's Creek, Kentucky. (E=eggs, R=rotifers, A=annelids, Cop=copepods, Clad=cladocerans, H=hydracarina, and D=dipterans).

Life Stage	Total length category (mm)	N	Mean TL	Mean gut											
				length (mm)	Gut:TL	E	R	A	Cop	Clad	H	D			
Pro-larvae		0													
Meso-larvae	6.00- 8.99	45	8.04	4.00	0.50:1	0.82	1.71	0.09	2.38	2.82	0	0			
	9.00-11.99	30	9.86	6.25	0.63:1	1.37	2.87	0.97	9.50	1.73	0	2.10			
	6.00-11.99	75	8.77	4.90	0.56:1	1.04	2.17	0.44	5.23	2.38	0	0.37			
Meta-larvae	8.00-10.99	13	9.54	5.69	0.60:1	0.23	1.54	0.08	4.61	3.00	0	1.77			
	11.00-13.99	17	12.37	8.48	0.68:1	1.29	0.18	0.06	5.53	1.82	0.06	5.35			
	14.00-16.99	14	15.52	9.23	0.59:1	1.93	0.21	0.43	9.07	3.14	0.14	4.57			
	17.00-19.99	14	18.32	13.21	0.72:1	2.50	0	0.71	11.28	1.36	0.21	8.71			
	20.00-23.00	18	21.69	17.87	0.82:1	1.56	0	0.11	17.44	1.61	0	4.83			
	8.00-23.00	76	15.77	11.24	0.71:1	1.05	0.34	0.26	9.90	2.13	0.08	5.09			

#### LITERATURE CITED

- Applegate, R. L. 1981. Food selection of muskellunge fry. *Prog. Fish-Cult.* 43:136-139.
- Bell, D. E. and R. D. Hoyt. 1980. Temporal and spatial abundance and diversity of fishes in a Kentucky stream. *Trans. Ky. Acad. Sci.* 41(1-2):35-44.
- Brocksen, R. W., G. E. Davis, and C. E. Warren. 1968. Competition, food consumption, and production of sculpins and trout in laboratory stream communities. *J. Wildl. Manage.* 32:51-75.
- Brooks, J. L. 1967. The effects of prey size selection by lake planktivores. *Sys. Zool.* 17:273-291.
- Brooks, J. L. and S. I. Dodson. 1965. Predation, body size, and composition of plankton. *Science* 150:28-35.
- Bulkley, R. V., V. L. Spykermann, and L. E. Inmon. 1976. Food of the pelagic young of walleyes and five cohabiting fish species in Clear Lake, Iowa. *Trans. Am. Fish. Soc.* 105:77-83.
- Chew, R. L. 1974. Early life history of the Florida largemouth bass. Florida Game and Freshwater Fisheries Commission Fishery Bulletin #7, 76 pp.
- Clark, A. L. and W. D. Pearson. 1979. Early piscivory in larvae of the freshwater drum, *Aplodinotus grunniens*. Pages 32-59 in Proceedings of a Workshop on Freshwater Larval Fishes. T. V. A. Division of Forestry, Fisheries and Wildlife Development, Norris, Tn.
- Confer, J. L., G. L. Howick, M. H. Corzette, S. L. Dramer, S. Fitzgibbon,

- and R. Landesberg. 1978. Visual predation by planktivores. *Oikos* 31:27-37.
- Echelle, A. A. 1968. Food habits of young-of-year longnose gar in Lake Texoma, Oklahoma. *Southwest. Nat.* 13(1):45-50.
- Eddy, S. and A. C. Hodson. 1970. Taxonomic keys to the common animals of the north central states. Burgess Publishing Co., Minneapolis, Minn. 162 pp.
- Elliott, J. M. and J. Corlett. 1971. The ecology of Morecambe Bay: invertebrate drift into and from the river Leven. *J. Appl. Ecol.* 9:195-205.
- Elton, C. 1946. Competition and structure of ecological communities. *J. Anim. Ecol.* 15:54-68.
- Floyd, K. B. 1983. Chronology of appearance and habitat partitioning by stream larval fishes. Unpublished Masters Thesis, Western Kentucky University, Bowling Green, Ky.
- Galbraith, M. G., Jr. 1967. Size-selection predation on *Daphnia* by rainbow trout and yellow perch. *Trans. Am. Fish. Soc.* 96:1-10.
- George, E. I. and W. F. Hadley. 1979. Partitioning between rock bass (*Ambloplites rupestris*) and smallmouth bass (*Micropterus dolomieu*) young of the year. *Trans. Am. Fish. Soc.* 108:253-261.
- Hansen, M. and D. H. Wahl. 1981. Selection of small *Daphnia pulex* by yellow perch fry in Oneida Lake, New York. *Trans. Am. Fish. Soc.* 110:64-71.
- Hartman, G. F. 1958. Mouth size and food size in young rainbow trout, *Salmo gairdneri*. *Copeia* 1958:233-234.



- Heins, D. C. and G. H. Clemmer. 1975. Ecology, foods and feeding of the longnose shiner, *Notropis longirostris* (Hay), in Mississippi. Am. Midl. Nat. 94:284-295.
- Hickling, C. F. 1966. On the feeding process in the white amur, *Ctenopharyngodon idella*. J. Zool. (Lond.). 148:408-419.
- Hogue, Jr., J. J., R. Wallus, and L. K. Kay. 1976. Preliminary guide to the identification of larval fishes in the Tennessee River. T. V. A. Division for Fishery and Wildlife Development, Norris, Tn. Tech. Note B-19, 66 pp.
- Hudson, P. L. and B. C. Cowell. 1966. Distribution and abundance of phytoplankton and rotifers in a main stem Missouri river reservoir. Proceedings of the South Dakota Academy of Science. 45:85-106.
- Hynes, H. B. N. 1970. The ecology of running waters. University of Toronto Press, Toronto, 555 pp.
- Ivlev, V. S. 1961. Experimental ecology of feeding of fishes (Translated by Douglass Scott). Yale University Press, New Haven, Pa., 302 pp.
- Kramer, R. H. and L. L. Smith, Jr. 1960. Formation of year classes in largemouth bass. Trans. Am. Fish. Soc. 91:29-41.
- Lagler, K. F., J. E. Bardoch, R. R. Miller, and D. R. Passino. 1977. Ichthyology. John Wiley & Sons, New York, N. Y., 506 pp.
- Larkin, P. A. 1956. Interspecific competition and population control in freshwater fish. J. Fish. Res. Board Can. 13:327-342.
- Mathur, D. 1977. Food habits and competitive relationships of the bandfin shiner in Halawakee Creek, Alabama. Am. Midl. Nat. 97:89-100.

- May, E. B. and C. R. Gasaway. 1967. A preliminary key to the identification of larval fishes of Oklahoma, with particular reference to Canton Reservoir, including a selected bibliography. Oklahoma Fish Research Laboratory Bulletin #5, Norman, Ok., 33 pp.
- Mendelson, J. 1975. Feeding relationships among species of *Notropis* (Pisces: Cyprinidae) in a Wisconsin stream. Ecol. Monogr. 45:199-230.
- Moore, J. W. and I. A. Moore. 1976. The basis of food selection in flounders, *Platichthys flesus* (L.), in the Severn Estuary. J. Fish Biol. 9:139-156.
- Nilsson, N. A. 1965. Food segregation between salmonid species in North Sweden. Rep. Inst. Freshwater Res. Drottningholm 46:58-78.
- Northcote, T. G. 1954. Observations on the comparative ecology of two species of fish, *Cottus asper* and *Cottus rhotheus* in British Columbia. Copeia 1954:25-28.
- Overmann, G. J., R. D. Hoyt, and G. A. Kindschi. 1980. The larval life history of the crappies (*Pomoxis*) spp. Trans. Ky. Acad. Sci. 41:105-115.
- Pennak, R. W. 1949. Annual limnological cycles in some Colorado lakes. Ecol. Monogr. 19:234-267.
- \_\_\_\_\_. 1978. Fresh-water invertebrates of the United States, 2nd ed. John W. Wiley & Sons, New York, N. Y., 803 pp.
- Peterson, A. 1973. Larvae of insects. Part II. Edwards Brothers, Inc. Ann Arbor, Mich., 416 pp.
- Pflieger, W. L. 1975. The fishes of Missouri. Missouri Department of Conservation, 343 pp.

- Scalet, C. G. 1972. Food habits of the orange belly darter, *Etheostoma radiosum cyanorum* (Osteichthes: Percidae). Am. Midl. Nat. 87:515-522.
- Selgeby, J. H. 1975. Life histories and abundance of crustacean zooplankton in the outlet of Lake Superior, 1971-72. J. Fish. Res. Board Can. 34:461-470.
- Siefert, R. E. 1968. Reproductive behavior, incubation and mortality of eggs, and postlarval food selection in the white crappie. Trans. Am. Fish. Soc. 97(3):252-259.
- \_\_\_\_\_. 1972. First food of larval yellow perch, white sucker, bluegill, emerald shiner, and rainbow smelt. Trans. Am. Fish. Soc. 101(2): 219-225.
- Silvells, H. C. 1949. Food studies of black crappie fry. Tex. J. Sci. 1:38-40.
- Snyder, D. E. 1976. Terminologies for intervals of larval fish development. Pages 41-58 in J. Boreman, ed. Great Lakes Fish Egg and Larval Identification. Office of Biological Services. U. S. Fish and Wildlife Service.
- Wallace, D. C. 1976. Feeding behavior and development, season and diel changes in the food of the silverjaw minnow, *Ericymba buccata* Cope. Am. Midl. Nat. 95:361-376.
- Ware, D. M. 1971. Predation by rainbow trout: the effect of experience. J. Fish. Res. Board Can. 28:1847-1852.
- Watkins, C., J. V. Shireman, R. W. Rottmann, and D. E. Colle. 1981. Food habits of fingerling grass carp. Prog. Fish-Cult. 43(2):95-97.

- Werner, E. E. 1979. Niche partitioning by food size in fish communities. Pages 331-322 in H. Clepper, ed. Predator-prey systems in fisheries management. Sport Fishing Institute, Washington, D. C.
- White, D. S. and K. H. Haag. 1977. Foods and feeding habits of the spotted sucker, *Minytrema melanops* (Rafinesque). Am. Midl. Nat. 98: 137-146.
- Winner, J. M. 1975. Zooplankton. Pages 155-169 in B. A. Whitton, ed. River ecology. University of California Press, Los Angeles, Ca.
- Wong, B. and F. J. Ward. 1972. Size selection of *Daphnia pulicaria* by yellow perch (*Perca flavescens*) fry in West Blue Lake, Manitoba. J. Fish. Res. Board Can. 29:1761-1764.
- Wynes, D. L. and T. E. Wissing. 1982. Resource sharing among darters in an Ohio stream. Am. Midl. Nat. 107:294-307.
- Zaret, T. M. and A. S. Rand. 1971. Competition in tropical stream fishes: support for the competitive exclusion principle. Ecology 52:336-342.