

## Mid Atlantic Regional Chapter of the American College of Sports Medicine



Annual Scientific Meeting, November 1<sup>st</sup> – 2<sup>nd</sup>, 2019

Conference Proceedings
International Journal of Exercise Science, Volume 9, Issue 8

## Effects of Capsaicin on the Hemodynamic Responses to Handgrip Exercise: Potential Influence of Race

Brian Lora, Tawn Tomasi, Meaghan Lynch, Gaia Giuriato, Jack Schickler, Emma Finegan, Emma Basso, Robert Restaino, and Stephen J. Ives. Skidmore College, Saratoga Springs, NY

Previous work in vitro suggests that capsaicin, the spicy ingredient in peppers, attenuates the vascular response to sympathetic activity ("sympatholysis"), likely mediated through vascular transient receptor potential vanilloid type 1 (TRPV1), improving vascular function, though translational studies are needed. Further, given racial disparities in cardiovascular health, exploring the potential physiological mechanisms underpinning these differences is warranted. Purpose: To determine the impact of acute capsaicin ingestion on the central hemodynamic responses to sympathetic stimuli, such as lower body negative pressure (LBNP) at rest. **Methods:** In a single-blind crossover design, 23 young healthy black (n=13) and white males (n=10) were given placebo (fiber) or capsaicin (Chile pepper) capsules. Stroke volume (SV), cardiac output (CO), mean arterial pressure (MAP), and heart rate (HR), were measured continuously using a Finometer, at rest and during LBNP (-20 mmHg). Results: Baseline SV was significantly different between black and white males (black male capsaicin: 81±15 vs white male capsaicin: 98±17, p=0.032). Baseline CO in the capsaicin trial was trending towards a significant difference (black male capsaicin: 4.6±0.8 vs white male capsaicin: 5.4±0.9, p=0.076). The LBNPinduced changes at rest appeared larger for black males (SV -18±4ml vs. -8±11ml and MAP 4.6±6 vs. 1.3±5mmHg, black vs. white), which were reversed with acute capsaicin (SV -10.7±13 vs. -9±16ml and MAP -2±9 vs. 1.2±8mmHg, black vs. white). Conclusion: Capsaicin, and assumed activation of TRPV1 receptors in afferent neurons, seems to affect the hemodynamic response to LBNP, in a race-specific manner. However, further work is needed to determine the mechanisms for the differences in hemodynamics at rest and in response to LBNP and acute capsaicin between races.